

to evaluate the ability of fusion proteins of the invention to act on hematopoietic stem cells in the context of the extracellular matrix (ECM) induced signal.

Cells respond to the regulatory factors in the context of signal(s) received from the surrounding microenvironment. For example, fibroblasts, and endothelial and epithelial stem cells fail to replicate in the absence of signals from the ECM. Hematopoietic stem cells can undergo self-renewal in the bone marrow, but not in *in vitro* suspension culture. The ability of stem cells to undergo self-renewal *in vitro* is dependent upon their interaction with the stromal cells and the ECM protein fibronectin (fn). Adhesion of cells to fn is mediated by the  $\alpha_5\beta_1$  and  $\alpha_4\beta_1$  integrin receptors, which are expressed by human and mouse hematopoietic stem cells. The factor(s) which integrate with the ECM environment and are responsible for stimulating stem cell self-renewal have not yet been identified. Discovery of such factors should be of great interest in gene therapy and bone marrow transplant applications

Briefly, polystyrene, non tissue culture treated, 96-well plates are coated with fn fragment at a coating concentration of 0.2  $\mu\text{g}/\text{cm}^2$ . Mouse bone marrow cells are plated (1,000 cells/well) in 0.2 ml of serum-free medium. Cells cultured in the presence of IL-3 (5 ng/ml) + SCF (50 ng/ml) would serve as the positive control, conditions under which little self-renewal but pronounced differentiation of the stem cells is to be expected. Albumin fusion proteins of the invention are tested with appropriate negative controls in the presence and absence of SCF(5.0 ng/ml), where volume of the administered composition containing the albumin fusion protein of the invention represents 10% of the total assay volume. The plated cells are then allowed to grow by incubating in a low oxygen environment (5%  $\text{CO}_2$ , 7%  $\text{O}_2$ , and 88%  $\text{N}_2$ ) tissue culture incubator for 7 days. The number of proliferating cells within the wells is then quantitated by measuring thymidine incorporation into cellular DNA. Verification of the positive hits in the assay will require phenotypic characterization of the cells, which can be accomplished by scaling up of the culture system and using appropriate antibody reagents against cell surface antigens and FACScan.

One skilled in the art could easily modify the exemplified studies to test the activity of albumin fusion proteins and polynucleotides of the invention (e.g., gene therapy).

If a particular fusion protein of the present invention is found to be a stimulator of hematopoietic progenitors, the fusion protein and polynucleotides corresponding to the fusion protein may be useful for example, in the diagnosis and treatment of disorders affecting the immune system and hematopoiesis. Representative uses are described in the "Immune Activity" and "Infectious Disease" sections above, and elsewhere herein. The fusion protein may also be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types.

Additionally, the albumin fusion proteins of the invention and polynucleotides encoding albumin fusion proteins of the invention, may also be employed to inhibit the

proliferation and differentiation of hematopoietic cells and therefore may be employed to protect bone marrow stem cells from chemotherapeutic agents during chemotherapy. This antiproliferative effect may allow administration of higher doses of chemotherapeutic agents and, therefore, more effective chemotherapeutic treatment.

5        Moreover, fusion proteins of the invention and polynucleotides encoding albumin fusion proteins of the invention may also be useful for the treatment and diagnosis of hematopoietic related disorders such as, anemia, pancytopenia, leukopenia, thrombocytopenia or leukemia, since stromal cells are important in the production of cells of hematopoietic lineages. The uses include bone marrow cell ex-vivo culture, bone marrow transplantation,  
10       bone marrow reconstitution, radiotherapy or chemotherapy of neoplasia.

#### Example 39: Human Dermal Fibroblast and Aortic Smooth Muscle Cell Proliferation

      An albumin fusion protein of the invention is added to cultures of normal human dermal fibroblasts (NHDF) and human aortic smooth muscle cells (AoSMC) and two co-  
15       assays are performed with each sample. The first assay examines the effect of the fusion protein on the proliferation of normal human dermal fibroblasts (NHDF) or aortic smooth muscle cells (AoSMC). Aberrant growth of fibroblasts or smooth muscle cells is a part of several pathological processes, including fibrosis, and restenosis. The second assay examines IL6 production by both NHDF and SMC. IL6 production is an indication of  
20       functional activation. Activated cells will have increased production of a number of cytokines and other factors, which can result in a proinflammatory or immunomodulatory outcome. Assays are run with and without co-TNF $\alpha$  stimulation, in order to check for costimulatory or inhibitory activity.

      Briefly, on day 1, 96-well black plates are set up with 1000 cells/well (NHDF) or  
25       2000 cells/well (AoSMC) in 100  $\mu$ l culture media. NHDF culture media contains: Clonetics FB basal media, 1mg/ml hFGF, 5mg/ml insulin, 50mg/ml gentamycin, 2%FBS, while AoSMC culture media contains Clonetics SM basal media, 0.5  $\mu$ g/ml hEGF, 5mg/ml insulin, 1 $\mu$ g/ml hFGF, 50mg/ml gentamycin, 50  $\mu$ g/ml Amphotericin B, 5%FBS. After incubation at 37°C for at least 4-5 hours culture media is aspirated and replaced with growth arrest media.  
30       Growth arrest media for NHDF contains fibroblast basal media, 50mg/ml gentamycin, 2% FBS, while growth arrest media for AoSMC contains SM basal media, 50mg/ml gentamycin, 50 $\mu$ g/ml Amphotericin B, 0.4% FBS. Incubate at 37 °C until day 2.

      On day 2, serial dilutions and templates of an albumin fusion protein of the invention are designed such that they always include media controls and known-protein controls. For  
35       both stimulation and inhibition experiments, proteins are diluted in growth arrest media. For inhibition experiments, TNF $\alpha$  is added to a final concentration of 2ng/ml (NHDF) or 5ng/ml

(AoSMC). Add 1/3 vol media containing controls or an albumin fusion protein of the invention and incubate at 37 degrees C/5% CO<sub>2</sub> until day 5.

Transfer 60  $\mu$ l from each well to another labeled 96-well plate, cover with a plate-sealer, and store at 4 degrees C until Day 6 (for IL6 ELISA). To the remaining 100  $\mu$ l in the cell culture plate, aseptically add Alamar Blue in an amount equal to 10% of the culture volume (10  $\mu$ l). Return plates to incubator for 3 to 4 hours. Then measure fluorescence with excitation at 530nm and emission at 590nm using the CytoFluor. This yields the growth stimulation/inhibition data.

On day 5, the IL6 ELISA is performed by coating a 96 well plate with 50-100  $\mu$ l/well of Anti-Human IL6 Monoclonal antibody diluted in PBS, pH 7.4, incubate ON at room temperature.

On day 6, empty the plates into the sink and blot on paper towels. Prepare Assay Buffer containing PBS with 4% BSA. Block the plates with 200  $\mu$ l/well of Pierce Super Block blocking buffer in PBS for 1-2 hr and then wash plates with wash buffer (PBS, 0.05% Tween-20). Blot plates on paper towels. Then add 50  $\mu$ l/well of diluted Anti-Human IL-6 Monoclonal, Biotin-labeled antibody at 0.50 mg/ml. Make dilutions of IL-6 stock in media (30, 10, 3, 1, 0.3, 0 ng/ml). Add duplicate samples to top row of plate. Cover the plates and incubate for 2 hours at RT on shaker.

Plates are washed with wash buffer and blotted on paper towels. Dilute EU-labeled Streptavidin 1:1000 in Assay buffer, and add 100  $\mu$ l/well. Cover the plate and incubate 1 h at RT. Plates are again washed with wash buffer and blotted on paper towels.

Add 100  $\mu$ l/well of Enhancement Solution. Shake for 5 minutes. Read the plate on the Wallac DELFIA Fluorometer. Readings from triplicate samples in each assay were tabulated and averaged.

A positive result in this assay suggests AoSMC cell proliferation and that the albumin fusion protein may be involved in dermal fibroblast proliferation and/or smooth muscle cell proliferation. A positive result also suggests many potential uses of the fusion protein and polynucleotides encoding the albumin fusion protein. For example, inflammation and immune responses, wound healing, and angiogenesis, as detailed throughout this specification. Particularly, fusion proteins may be used in wound healing and dermal regeneration, as well as the promotion of vasculogenesis, both of the blood vessels and lymphatics. The growth of vessels can be used in the treatment of, for example, cardiovascular diseases. Additionally, fusion proteins showing antagonistic activity in this assay may be useful in treating diseases, disorders, and/or conditions which involve angiogenesis by acting as an anti-vascular agent (e.g., anti-angiogenesis). These diseases, disorders, and/or conditions are known in the art and/or are described herein, such as, for

example, malignancies, solid tumors, benign tumors, for example hemangiomas, acoustic neuromas, neurofibromas, trachomas, and pyogenic granulomas; arteriosclerotic plaques; ocular angiogenic diseases, for example, diabetic retinopathy, retinopathy of prematurity, macular degeneration, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, rubeosis, retinoblastoma, uveitis and Pterygia (abnormal blood vessel growth) of the eye; rheumatoid arthritis; psoriasis; delayed wound healing; endometriosis; vasculogenesis; granulations; hypertrophic scars (keloids); nonunion fractures; scleroderma; trachoma; vascular adhesions; myocardial angiogenesis; coronary collaterals; cerebral collaterals; arteriovenous malformations; ischemic limb angiogenesis; Osler-Webber Syndrome; plaque neovascularization; telangiectasia; hemophilic joints; angiofibroma; fibromuscular dysplasia; wound granulation; Crohn's disease; and atherosclerosis. Moreover, albumin fusion proteins that act as antagonists in this assay may be useful in treating anti-hyperproliferative diseases and/or anti-inflammatory known in the art and/or described herein.

*Example 40: Cellular Adhesion Molecule (CAM) Expression on Endothelial Cells*

The recruitment of lymphocytes to areas of inflammation and angiogenesis involves specific receptor-ligand interactions between cell surface adhesion molecules (CAMs) on lymphocytes and the vascular endothelium. The adhesion process, in both normal and pathological settings, follows a multi-step cascade that involves intercellular adhesion molecule-1 (ICAM-1), vascular cell adhesion molecule-1 (VCAM-1), and endothelial leukocyte adhesion molecule-1 (E-selectin) expression on endothelial cells (EC). The expression of these molecules and others on the vascular endothelium determines the efficiency with which leukocytes may adhere to the local vasculature and extravasate into the local tissue during the development of an inflammatory response. The local concentration of cytokines and growth factor participate in the modulation of the expression of these CAMs.

Briefly, endothelial cells (e.g., Human Umbilical Vein Endothelial cells (HUVECs)) are grown in a standard 96 well plate to confluence, growth medium is removed from the cells and replaced with 100  $\mu$ l of 199 Medium (10% fetal bovine serum (FBS)). Samples for testing (containing an albumin fusion protein of the invention) and positive or negative controls are added to the plate in triplicate (in 10  $\mu$ l volumes). Plates are then incubated at 37°C for either 5 h (selectin and integrin expression) or 24 h (integrin expression only). Plates are aspirated to remove medium and 100  $\mu$ l of 0.1% paraformaldehyde-PBS(with Ca++ and Mg++) is added to each well. Plates are held at 4°C for 30 min. Fixative is removed from the wells and wells are washed 1X with PBS(+Ca,Mg) + 0.5% BSA and drained. 10  $\mu$ l of diluted primary antibody is added to the test and control wells. Anti-ICAM-1-Biotin, Anti-VCAM-1-Biotin and Anti-E-selectin-Biotin are used at a concentration of 10



$\mu\text{g/ml}$  (1:10 dilution of 0.1 mg/ml stock antibody). Cells are incubated at 37°C for 30 min. in a humidified environment. Wells are washed three times with PBS(+Ca,Mg) + 0.5% BSA. 20  $\mu\text{l}$  of diluted ExtrAvidin-Alkaline Phosphatase (1:5,000 dilution, referred to herein as the working dilution) are added to each well and incubated at 37°C for 30 min. Wells are washed  
5 three times with PBS(+Ca,Mg)+0.5% BSA. Dissolve 1 tablet of p-Nitrophenol Phosphate pNPP per 5 ml of glycine buffer (pH 10.4). 100  $\mu\text{l}$  of pNPP substrate in glycine buffer is added to each test well. Standard wells in triplicate are prepared from the working dilution of the ExtrAvidin-Alkaline Phosphatase in glycine buffer: 1:5,000 ( $10^0$ ) >  $10^{-0.5}$  >  $10^{-1}$  >  $10^{-1.5}$ . 5  
10  $\mu\text{l}$  of each dilution is added to triplicate wells and the resulting AP content in each well is 5.50 ng, 1.74 ng, 0.55 ng, 0.18 ng. 100  $\mu\text{l}$  of pNPP reagent is then added to each of the standard wells. The plate is incubated at 37°C for 4h. A volume of 50  $\mu\text{l}$  of 3M NaOH is added to all wells. The plate is read on a plate reader at 405 nm using the background subtraction option on blank wells filled with glycine buffer only. Additionally, the template is set up to indicate the concentration of AP-conjugate in each standard well [ 5.50 ng; 1.74 ng;  
15 0.55 ng; 0.18 ng]. Results are indicated as amount of bound AP-conjugate in each sample.

#### ***Example 41: Alamar Blue Endothelial Cells Proliferation Assay***

This assay may be used to quantitatively determine protein mediated inhibition of bFGF-induced proliferation of Bovine Lymphatic Endothelial Cells (LECs), Bovine Aortic  
20 Endothelial Cells (BAECs) or Human Microvascular Uterine Myometrial Cells (UTMECs). This assay incorporates a fluorometric growth indicator based on detection of metabolic activity. A standard Alamar Blue Proliferation Assay is prepared in EGM-2MV with 10 ng /ml of bFGF added as a source of endothelial cell stimulation. This assay may be used with a variety of endothelial cells with slight changes in growth medium and cell concentration.  
25 Dilutions of protein batches to be tested are diluted as appropriate. Serum-free medium (GIBCO SFM) without bFGF is used as a non-stimulated control and Angiostatin or TSP-1 are included as a known inhibitory controls.

Briefly, LEC, BAECs or UTMECs are seeded in growth media at a density of 5000 to 2000 cells/well in a 96 well plate and placed at 37 degreesC overnight. After the overnight  
30 incubation of the cells, the growth media is removed and replaced with GIBCO EC-SFM. The cells are treated with the appropriate dilutions of an albumin fusion protein of the invention or control protein sample(s) (prepared in SFM ) in triplicate wells with additional bFGF to a concentration of 10 ng/ ml. Once the cells have been treated with the samples, the plate(s) is/are placed back in the 37° C incubator for three days. After three days 10 ml of  
35 stock alamar blue (Biosource Cat# DAL1100) is added to each well and the plate(s) is/are placed back in the 37°C incubator for four hours. The plate(s) are then read at 530nm excitation and 590nm emission using the CytoFluor fluorescence reader. Direct output is

recorded in relative fluorescence units.

Alamar blue is an oxidation-reduction indicator that both fluoresces and changes color in response to chemical reduction of growth medium resulting from cell growth. As cells grow in culture, innate metabolic activity results in a chemical reduction of the immediate surrounding environment. Reduction related to growth causes the indicator to change from oxidized (non-fluorescent blue) form to reduced (fluorescent red) form (i.e., stimulated proliferation will produce a stronger signal and inhibited proliferation will produce a weaker signal and the total signal is proportional to the total number of cells as well as their metabolic activity). The background level of activity is observed with the starvation medium alone.

This is compared to the output observed from the positive control samples (bFGF in growth medium) and protein dilutions.

#### *Example 42: Detection of Inhibition of a Mixed Lymphocyte Reaction*

This assay can be used to detect and evaluate inhibition of a Mixed Lymphocyte Reaction (MLR) by fusion proteins of the invention. Inhibition of a MLR may be due to a direct effect on cell proliferation and viability, modulation of costimulatory molecules on interacting cells, modulation of adhesiveness between lymphocytes and accessory cells, or modulation of cytokine production by accessory cells. Multiple cells may be targeted by the albumin fusion proteins that inhibit MLR since the peripheral blood mononuclear fraction used in this assay includes T, B and natural killer lymphocytes, as well as monocytes and dendritic cells.

Albumin fusion proteins of the invention found to inhibit the MLR may find application in diseases associated with lymphocyte and monocyte activation or proliferation. These include, but are not limited to, diseases such as asthma, arthritis, diabetes, inflammatory skin conditions, psoriasis, eczema, systemic lupus erythematosus, multiple sclerosis, glomerulonephritis, inflammatory bowel disease, crohn's disease, ulcerative colitis, arteriosclerosis, cirrhosis, graft vs. host disease, host vs. graft disease, hepatitis, leukemia and lymphoma.

Briefly, PBMCs from human donors are purified by density gradient centrifugation using Lymphocyte Separation Medium (LSM<sup>®</sup>, density 1.0770 g/ml, Organon Teknika Corporation, West Chester, PA). PBMCs from two donors are adjusted to  $2 \times 10^6$  cells/ml in RPMI-1640 (Life Technologies, Grand Island, NY) supplemented with 10% FCS and 2 mM glutamine. PBMCs from a third donor is adjusted to  $2 \times 10^5$  cells/ml. Fifty microliters of PBMCs from each donor is added to wells of a 96-well round bottom microtiter plate.

Dilutions of the fusion protein test material (50  $\mu$ l) is added in triplicate to microtiter wells. Test samples (of the protein of interest) are added for final dilution of 1:4; rhuIL-2 (R&D Systems, Minneapolis, MN, catalog number 202-IL) is added to a final concentration of 1

5  $\mu\text{g/ml}$ ; anti-CD4 mAb (R&D Systems, clone 34930.11, catalog number MAB379) is added to a final concentration of 10  $\mu\text{g/ml}$ . Cells are cultured for 7-8 days at 37°C in 5% CO<sub>2</sub>, and 1  $\mu\text{C}$  of [<sup>3</sup>H] thymidine is added to wells for the last 16 hrs of culture. Cells are harvested and thymidine incorporation determined using a Packard TopCount. Data is expressed as the mean and standard deviation of triplicate determinations.

Samples of the fusion protein of interest are screened in separate experiments and compared to the negative control treatment, anti-CD4 mAb, which inhibits proliferation of lymphocytes and the positive control treatment, IL-2 (either as recombinant material or supernatant), which enhances proliferation of lymphocytes.

#### Example 43: Assays for Protease Activity

The following assay may be used to assess protease activity of an albumin fusion protein of the invention.

15 Gelatin and casein zymography are performed essentially as described (Heusen et al., *Anal. Biochem.*, 102:196-202 (1980); Wilson et al., *Journal of Urology*, 149:653-658 (1993)). Samples are run on 10% polyacrylamide/0.1% SDS gels containing 1% gelatin or casein, soaked in 2.5% triton at room temperature for 1 hour, and in 0.1M glycine, pH 8.3 at 37°C 5 to 16 hours. After staining in amido black areas of proteolysis appear as clear areas against the blue-black background. Trypsin (Sigma T8642) is used as a positive control.

20 Protease activity is also determined by monitoring the cleavage of n-a-benzoyl-L-arginine ethyl ester (BAEE) (Sigma B-4500. Reactions are set up in (25mM NaPO<sub>4</sub>, 1mM EDTA, and 1mM BAEE), pH 7.5. Samples are added and the change in adsorbance at 260nm is monitored on the Beckman DU-6 spectrophotometer in the time-drive mode. Trypsin is used as a positive control.

25 Additional assays based upon the release of acid-soluble peptides from casein or hemoglobin measured as adsorbance at 280 nm or colorimetrically using the Folin method are performed as described in Bergmeyer, et al., *Methods of Enzymatic Analysis*, 5 (1984). Other assays involve the solubilization of chromogenic substrates (Ward, *Applied Science*, 251-317 (1983)).

#### Example 44: Identifying Serine Protease Substrate Specificity

30 Methods known in the art or described herein may be used to determine the substrate specificity of the albumin fusion proteins of the present invention having serine protease activity. A preferred method of determining substrate specificity is by the use of positional scanning synthetic combinatorial libraries as described in GB 2 324 529 (incorporated herein in its entirety).

**Example 45: Ligand Binding Assays**

The following assay may be used to assess ligand binding activity of an albumin fusion protein of the invention.

Ligand binding assays provide a direct method for ascertaining receptor pharmacology and are adaptable to a high throughput format. The purified ligand for an albumin fusion protein of the invention is radiolabeled to high specific activity (50-2000 Ci/mmol) for binding studies. A determination is then made that the process of radiolabeling does not diminish the activity of the ligand towards the fusion protein. Assay conditions for buffers, ions, pH and other modulators such as nucleotides are optimized to establish a workable signal to noise ratio for both membrane and whole cell polypeptide sources. For these assays, specific polypeptide binding is defined as total associated radioactivity minus the radioactivity measured in the presence of an excess of unlabeled competing ligand. Where possible, more than one competing ligand is used to define residual nonspecific binding.

**Example 46: Functional Assay in *Xenopus* Oocytes**

Capped RNA transcripts from linearized plasmid templates encoding an albumin fusion protein of the invention is synthesized in vitro with RNA polymerases in accordance with standard procedures. In vitro transcripts are suspended in water at a final concentration of 0.2 mg/ml. Ovarian lobes are removed from adult female toads, Stage V defolliculated oocytes are obtained, and RNA transcripts (10 ng/oocyte) are injected in a 50 nl bolus using a microinjection apparatus. Two electrode voltage clamps are used to measure the currents from individual *Xenopus* oocytes in response fusion protein and polypeptide agonist exposure. Recordings are made in Ca<sup>2+</sup> free Barth's medium at room temperature. The *Xenopus* system can be used to screen known ligands and tissue/cell extracts for activating ligands.

**Example 47: Microphysiometric Assays**

Activation of a wide variety of secondary messenger systems results in extrusion of small amounts of acid from a cell. The acid formed is largely as a result of the increased metabolic activity required to fuel the intracellular signaling process. The pH changes in the media surrounding the cell are very small but are detectable by the CYTOSENSOR microphysiometer (Molecular Devices Ltd., Menlo Park, Calif.). The CYTOSENSOR is thus capable of detecting the ability of an albumin fusion protein of the invention to activate secondary messengers that are coupled to an energy utilizing intracellular signaling pathway.

**Example 48: Extract/Cell Supernatant Screening**

A large number of mammalian receptors exist for which there remains, as yet, no cognate activating ligand (agonist). Thus, active ligands for these receptors may not be included within the ligands banks as identified to date. Accordingly, the albumin fusion proteins of the invention can also be functionally screened (using calcium, cAMP, microphysiometer, oocyte electrophysiology, etc., functional screens) against tissue extracts to identify natural ligands for the Therapeutic protein portion and/or albumin protein portion of an albumin fusion protein of the invention. Extracts that produce positive functional responses can be sequentially subfractionated until an activating ligand is isolated and identified.

#### *Example 49: ATP-binding assay*

The following assay may be used to assess ATP-binding activity of fusion proteins of the invention.

ATP-binding activity of an albumin fusion protein of the invention may be detected using the ATP-binding assay described in U.S. Patent 5,858,719, which is herein incorporated by reference in its entirety. Briefly, ATP-binding to an albumin fusion protein of the invention is measured via photoaffinity labeling with 8-azido-ATP in a competition assay. Reaction mixtures containing 1 mg/ml of ABC transport protein are incubated with varying concentrations of ATP, or the non-hydrolyzable ATP analog adenylyl-5'-imidodiphosphate for 10 minutes at 4°C. A mixture of 8-azido-ATP (Sigma Chem. Corp., St. Louis, MO.) plus 8-azido-ATP ( $^{32}\text{P}$ -ATP) (5 mCi/ $\mu\text{mol}$ , ICN, Irvine CA.) is added to a final concentration of 100  $\mu\text{M}$  and 0.5 ml aliquots are placed in the wells of a porcelain spot plate on ice. The plate is irradiated using a short wave 254 nm UV lamp at a distance of 2.5 cm from the plate for two one-minute intervals with a one-minute cooling interval in between. The reaction is stopped by addition of dithiothreitol to a final concentration of 2mM. The incubations are subjected to SDS-PAGE electrophoresis, dried, and autoradiographed. Protein bands corresponding to the albumin fusion proteins of the invention are excised, and the radioactivity quantified. A decrease in radioactivity with increasing ATP or adenylyl-5'-imidodiphosphate provides a measure of ATP affinity to the fusion protein.

#### *Example 50: Phosphorylation Assay*

In order to assay for phosphorylation activity of an albumin fusion protein of the invention, a phosphorylation assay as described in U.S. Patent 5,958,405 (which is herein incorporated by reference) is utilized. Briefly, phosphorylation activity may be measured by phosphorylation of a protein substrate using gamma-labeled  $^{32}\text{P}$ -ATP and quantitation of the incorporated radioactivity using a gamma radioisotope counter. The fusion protein of the invention is incubated with the protein substrate,  $^{32}\text{P}$ -ATP, and a kinase buffer. The  $^{32}\text{P}$

incorporated into the substrate is then separated from free  $^{32}\text{P}$ -ATP by electrophoresis, and the incorporated  $^{32}\text{P}$  is counted and compared to a negative control. Radioactivity counts above the negative control are indicative of phosphorylation activity of the fusion protein.

5           ***Example 51: Detection of Phosphorylation Activity (Activation) of an Albumin Fusion Protein of the Invention in the Presence of Polypeptide Ligands***

Methods known in the art or described herein may be used to determine the phosphorylation activity of an albumin fusion protein of the invention. A preferred method of  
10 determining phosphorylation activity is by the use of the tyrosine phosphorylation assay as described in US 5,817,471 (incorporated herein by reference).

***Example 52: Identification Of Signal Transduction Proteins That Interact With An albumin fusion protein Of The Present Invention***

15           Albumin fusion proteins of the invention may serve as research tools for the identification, characterization and purification of signal transduction pathway proteins or receptor proteins. Briefly, a labeled fusion protein of the invention is useful as a reagent for the purification of molecules with which it interacts. In one embodiment of affinity purification, an albumin fusion protein of the invention is covalently coupled to a  
20 chromatography column. Cell-free extract derived from putative target cells, such as carcinoma tissues, is passed over the column, and molecules with appropriate affinity bind to the albumin fusion protein. The protein complex is recovered from the column, dissociated, and the recovered molecule subjected to N-terminal protein sequencing. This amino acid sequence is then used to identify the captured molecule or to design degenerate  
25 oligonucleotide probes for cloning the relevant gene from an appropriate cDNA library.

***Example 53: IL-6 Bioassay***

A variety of assays are known in the art for testing the proliferative effects of an albumin fusion protein of the invention. For example, one such assay is the IL-6 Bioassay  
30 as described by Marz *et al.* (*Proc. Natl. Acad. Sci., U.S.A.*, 95:3251-56 (1998)), which is herein incorporated by reference). After 68 hrs. at 37°C, the number of viable cells is measured by adding the tetrazolium salt thiazolyl blue (MTT) and incubating for a further 4 hrs. at 37°C. B9 cells are lysed by SDS and optical density is measured at 570 nm. Controls containing IL-6 (positive) and no cytokine (negative) are Briefly, IL-6 dependent B9 murine  
35 cells are washed three times in IL-6 free medium and plated at a concentration of 5,000 cells per well in 50  $\mu\text{l}$ , and 50  $\mu\text{l}$  of fusion protein of the invention is added. Enhanced proliferation in the test sample(s) (containing an albumin fusion protein of the invention)

relative to the negative control is indicative of proliferative effects mediated by the fusion protein.

**Example 54: Support of Chicken Embryo Neuron Survival**

To test whether sympathetic neuronal cell viability is supported by an albumin fusion protein of the invention, the chicken embryo neuronal survival assay of Senaldi *et al* may be utilized (*Proc. Natl. Acad. Sci., U.S.A.*, 96:11458-63 (1998), which is herein incorporated by reference). Briefly, motor and sympathetic neurons are isolated from chicken embryos, resuspended in L15 medium (with 10% FCS, glucose, sodium selenite, progesterone, conalbumin, putrescine, and insulin; Life Technologies, Rockville, MD.) and Dulbecco's modified Eagles medium [with 10% FCS, glutamine, penicillin, and 25 mM Hepes buffer (pH 7.2); Life Technologies, Rockville, MD.], respectively, and incubated at 37°C in 5% CO<sub>2</sub> in the presence of different concentrations of the purified fusion protein of the invention, as well as a negative control lacking any cytokine. After 3 days, neuron survival is determined by evaluation of cellular morphology, and through the use of the colorimetric assay of Mosmann (Mosmann, T., *J. Immunol. Methods*, 65:55-63 (1983)). Enhanced neuronal cell viability as compared to the controls lacking cytokine is indicative of the ability of the albumin fusion protein to enhance the survival of neuronal cells.

**Example 55: Assay for Phosphatase Activity**

The following assay may be used to assess serine/threonine phosphatase (PTPase) activity of an albumin fusion protein of the invention.

In order to assay for serine/threonine phosphatase (PTPase) activity, assays can be utilized which are widely known to those skilled in the art. For example, the serine/threonine phosphatase (PSPase) activity of an albumin fusion protein of the invention may be measured using a PSPase assay kit from New England Biolabs, Inc. Myelin basic protein (MyBP), a substrate for PSPase, is phosphorylated on serine and threonine residues with cAMP-dependent Protein Kinase in the presence of [<sup>32</sup>P]ATP. Protein serine/threonine phosphatase activity is then determined by measuring the release of inorganic phosphate from <sup>32</sup>P-labeled MyBP.

**Example 56: Interaction of Serine/Threonine Phosphatases with other Proteins**

Fusion protein of the invention having serine/threonine phosphatase activity (e.g., as determined in Example 55) are useful, for example, as research tools for the identification, characterization and purification of additional interacting proteins or receptor proteins, or other signal transduction pathway proteins. Briefly, a labeled fusion protein of the invention is

useful as a reagent for the purification of molecules with which it interacts. In one embodiment of affinity purification, an albumin fusion protein of the invention is covalently coupled to a chromatography column. Cell-free extract derived from putative target cells, such as neural or liver cells, is passed over the column, and molecules with appropriate affinity bind to the fusion protein. The fusion protein -complex is recovered from the column, dissociated, and the recovered molecule subjected to N-terminal protein sequencing. This amino acid sequence is then used to identify the captured molecule or to design degenerate oligonucleotide probes for cloning the relevant gene from an appropriate cDNA library.

#### *Example 57: Assaying for Heparanase Activity*

There are numerous assays known in the art that may be employed to assay for heparanase activity of an albumin fusion protein of the invention. In one example, heparanase activity of an albumin fusion protein of the invention, is assayed as described by Vlodavsky et al., (Vlodavsky et al., Nat. Med., 5:793-802 (1999)). Briefly, cell lysates, conditioned media, intact cells ( $1 \times 10^6$  cells per 35-mm dish), cell culture supernatant, or purified fusion protein are incubated for 18 hrs at 37°C, pH 6.2-6.6, with  $^{35}\text{S}$ -labeled ECM or soluble ECM derived peak I proteoglycans. The incubation medium is centrifuged and the supernatant is analyzed by gel filtration on a Sepharose CL-6B column (0.9 x 30 cm). Fractions are eluted with PBS and their radioactivity is measured. Degradation fragments of heparan sulfate side chains are eluted from Sepharose 6B at  $0.5 < K_{av} < 0.8$  (peak II). Each experiment is done at least three times. Degradation fragments corresponding to "peak II," as described by Vlodavsky et al., is indicative of the activity of an albumin fusion protein of the invention in cleaving heparan sulfate.

#### *Example 58: Immobilization of biomolecules*

This example provides a method for the stabilization of an albumin fusion protein of the invention in non-host cell lipid bilayer constructs (see, e.g., Bieri et al., Nature Biotech 17:1105-1108 (1999), hereby incorporated by reference in its entirety herein) which can be adapted for the study of fusion proteins of the invention in the various functional assays described above. Briefly, carbohydrate-specific chemistry for biotinylation is used to confine a biotin tag to an albumin fusion protein of the invention, thus allowing uniform orientation upon immobilization. A 50uM solution of an albumin fusion protein of the invention in washed membranes is incubated with 20 mM NaIO<sub>4</sub> and 1.5 mg/ml (4mM) BACH or 2 mg/ml (7.5mM) biotin-hydrazide for 1 hr at room temperature (reaction volume, 150ul). Then the sample is dialyzed (Pierce Slidealizer Cassett, 10 kDa cutoff; Pierce Chemical Co., Rockford IL) at 4°C first for 5 h, exchanging the buffer after each hour, and finally for 12 h against 500 ml buffer R (0.15 M NaCl, 1 mM MgCl<sub>2</sub>, 10 mM sodium phosphate, pH7). Just



before addition into a cuvette, the sample is diluted 1:5 in buffer ROG50 (Buffer R supplemented with 50 mM octylglucoside).

*Example 59: Assays for Metalloproteinase Activity*

5 Metalloproteinases are peptide hydrolases which use metal ions, such as  $Zn^{2+}$ , as the catalytic mechanism. Metalloproteinase activity of an albumin fusion protein of the present invention can be assayed according to methods known in the art. The following exemplary methods are provided:

*Proteolysis of alpha-2-macroglobulin*

10 To confirm protease activity, a purified fusion protein of the invention is mixed with the substrate alpha-2-macroglobulin (0.2 unit/ml; Boehringer Mannheim, Germany) in 1x assay buffer (50 mM HEPES, pH 7.5, 0.2 M NaCl, 10 mM  $CaCl_2$ , 25  $\mu M$   $ZnCl_2$  and 0.05% Brij-35) and incubated at 37°C for 1-5 days. Trypsin is used as positive control. Negative controls contain only alpha-2-macroglobulin in assay buffer. The samples are collected and  
15 boiled in SDS-PAGE sample buffer containing 5% 2-mercaptoethanol for 5-min, then loaded onto 8% SDS-polyacrylamide gel. After electrophoresis the proteins are visualized by silver staining. Proteolysis is evident by the appearance of lower molecular weight bands as compared to the negative control.

20 *Inhibition of alpha-2-macroglobulin proteolysis by inhibitors of metalloproteinases*

Known metalloproteinase inhibitors (metal chelators (EDTA, EGTA, AND  $HgCl_2$ ), peptide metalloproteinase inhibitors (TIMP-1 and TIMP-2), and commercial small molecule MMP inhibitors) may also be used to characterize the proteolytic activity of an albumin fusion protein of the invention. Three synthetic MMP inhibitors that may be used are: MMP  
25 inhibitor I, [ $IC_{50}$  = 1.0  $\mu M$  against MMP-1 and MMP-8;  $IC_{50}$  = 30  $\mu M$  against MMP-9;  $IC_{50}$  = 150  $\mu M$  against MMP-3]; MMP-3 (stromelysin-1) inhibitor I [ $IC_{50}$  = 5  $\mu M$  against MMP-3], and MMP-3 inhibitor II [ $K_i$  = 130 nM against MMP-3]; inhibitors available through Calbiochem, catalog # 444250, 444218, and 444225, respectively). Briefly, different concentrations of the small molecule MMP inhibitors are mixed with a purified fusion protein  
30 of the invention (50 $\mu g$ /ml) in 22.9  $\mu l$  of 1x HEPES buffer (50 mM HEPES, pH 7.5, 0.2 M NaCl, 10 mM  $CaCl_2$ , 25  $\mu M$   $ZnCl_2$  and 0.05%Brij-35) and incubated at room temperature (24 °C) for 2-hr, then 7.1  $\mu l$  of substrate alpha-2-macroglobulin (0.2 unit/ml) is added and incubated at 37°C for 20-hr. The reactions are stopped by adding 4x sample buffer and boiled immediately for 5 minutes. After SDS-PAGE, the protein bands are visualized by silver stain.  
35

*Synthetic Fluorogenic Peptide Substrates Cleavage Assay*

The substrate specificity for fusion proteins of the invention with demonstrated metalloproteinase activity may be determined using techniques known in the art, such as using synthetic fluorogenic peptide substrates (purchased from BACHEM Bioscience Inc). Test substrates include, M-1985, M-2225, M-2105, M-2110, and M-2255. The first four are MMP substrates and the last one is a substrate of tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) converting enzyme (TACE). These substrates are preferably prepared in 1:1 dimethyl sulfoxide (DMSO) and water. The stock solutions are 50-500  $\mu$ M. Fluorescent assays are performed by using a Perkin Elmer LS 50B luminescence spectrometer equipped with a constant temperature water bath. The excitation  $\lambda$  is 328 nm and the emission  $\lambda$  is 393 nm. Briefly, the assay is carried out by incubating 176  $\mu$ l 1x HEPES buffer (0.2 M NaCl, 10 mM  $\text{CaCl}_2$ , 0.05% Brij-35 and 50 mM HEPES, pH 7.5) with 4  $\mu$ l of substrate solution (50  $\mu$ M) at 25  $^\circ$ C for 15 minutes, and then adding 20  $\mu$ l of a purified fusion protein of the invention into the assay cuvette. The final concentration of substrate is 1  $\mu$ M. Initial hydrolysis rates are monitored for 30-min.

It will be clear that the invention may be practiced otherwise than as particularly described in the foregoing description and examples. Numerous modifications and variations of the present invention are possible in light of the above teachings and, therefore, are within the scope of the appended claims.

The entire disclosure of each document cited (including patents, patent applications, patent publications, journal articles, abstracts, laboratory manuals, books, or other disclosures) as well as information available through Identifiers specific to databases such as GenBank, GeneSeq, or the CAS Registry, referred to in this application are herein incorporated by reference in their entirety. The specification and sequence listing of each of the following U.S. applications are herein incorporated by reference in their entirety: Application Nos. 60/229,358 filed on April 12, 2000; 60/199,384 filed on April 25, 2000 and 60/256,931 filed on December 21, 2000.

**INDICATIONS RELATING TO A DEPOSITED MICROORGANISM  
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 59, line 12.

**B. IDENTIFICATION OF DEPOSIT**

Further deposits are identified on an additional sheet ☒

Name of depositary institution: American Type Culture Collection

Address of depositary institution (including postal code and country)

10801 University Boulevard  
Manassas, Virginia 20110-2209  
United States of America

Date of deposit

11 April 2001

Accession Number

Unassigned

**C. ADDITIONAL INDICATIONS** (leave blank if not applicable)

This information is continued on an additional sheet ☐

**D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE** (if the indications are not for all designated States)

Europe

In respect of those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28(4) EPC).

Continued on additional sheets

**E. SEPARATE FURNISHING OF INDICATIONS** (leave blank if not applicable)

The indications listed below will be submitted to the international Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")

For receiving Office use only		For International Bureau use only	
<input checked="" type="checkbox"/> This sheet was received with the international application <b>Enora Rivera</b> <b>PET Operations - IAPD Team 1</b> Authorized officer: (703) 305-3678 (703) 305-3230 (FAX)		<input type="checkbox"/> This sheet was received by the International Bureau on:  Authorized officer:	

**ATCC Deposit No.: Unassigned**

## **CANADA**

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

## **NORWAY**

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

## **AUSTRALIA**

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

## **FINLAND**

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

## **UNITED KINGDOM**

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

**ATCC Deposit No.: Unassigned**

## **DENMARK**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

## **SWEDEN**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

## **NETHERLANDS**

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

# **INDICATIONS RELATING TO A DEPOSITED MICROORGANISM OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 58, line 12.

## **B. IDENTIFICATION OF DEPOSIT**

Further deposits are identified on an additional sheet ☒

Name of depositary institution: American Type Culture Collection

Address of depositary institution (including postal code and country)

10801 University Boulevard  
Manassas, Virginia 20110-2209  
United States of America

Date of deposit

11 April 2001

Accession Number

Unassigned

## **C. ADDITIONAL INDICATIONS** (leave blank if not applicable)

This information is continued on an additional sheet ☐

## **D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE** (if the indications are not for all designated States)

Europe

In respect of those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28(4) EPC).

Continued on additional sheets

## **E. SEPARATE FURNISHING OF INDICATIONS** (leave blank if not applicable)

The indications listed below will be submitted to the international Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")

For receiving Office use only	For International Bureau use only	
<input checked="" type="checkbox"/> This sheet was received with the international application <b>Enara Rivera</b> <b>PCT Operations - IAPD Team 1</b> Authorized officer (703) 305-3678 (703) 305-3230 (FAX)		<input type="checkbox"/> This sheet was received by the International Bureau on:  Authorized officer

**ATCC Deposit No.: Unassigned**

## **CANADA**

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

## **NORWAY**

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

## **AUSTRALIA**

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

## **FINLAND**

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

## **UNITED KINGDOM**

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

**ATCC Deposit No.: Unassigned**

## **DENMARK**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

## **SWEDEN**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

## **NETHERLANDS**

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.



**INDICATIONS RELATING TO A DEPOSITED MICROORGANISM  
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 59, line 12.

**B. IDENTIFICATION OF DEPOSIT**

Further deposits are identified on an additional sheet ☒

Name of depositary institution: American Type Culture Collection

Address of depositary institution (including postal code and country)

10801 University Boulevard  
Manassas, Virginia 20110-2209  
United States of America

Date of deposit

11 April 2001

Accession Number

Unassigned

**C. ADDITIONAL INDICATIONS** (leave blank if not applicable)

This information is continued on an additional sheet ☐

**D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE** (if the indications are not for all designated States)

Europe

In respect of those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28(4) EPC).

Continued on additional sheets

**E. SEPARATE FURNISHING OF INDICATIONS** (leave blank if not applicable)

The indications listed below will be submitted to the international Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")

For receiving Office use only		For International Bureau use only	
<input checked="" type="checkbox"/> This sheet was received with the international application <b>Elmora Rivera</b> <b>PCT Operations - IAPD Team 1</b> Authorized officer: <b>(703) 305-3678 (703) 305-3230 (FAX)</b>		<input type="checkbox"/> This sheet was received by the International Bureau on: Authorized officer:	

**ATCC Deposit No.: Unassigned****CANADA**

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

**NORWAY**

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

**AUSTRALIA**

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

**FINLAND**

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

**UNITED KINGDOM**

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

**ATCC Deposit No.: Unassigned**

## **DENMARK**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

## **SWEDEN**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

## **NETHERLANDS**

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

**INDICATIONS RELATING TO A DEPOSITED MICROORGANISM  
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 59, line 12.

**B. IDENTIFICATION OF DEPOSIT**

Further deposits are identified on an additional sheet ☒

Name of depositary institution: American Type Culture Collection

Address of depositary institution (including postal code and country)

10801 University Boulevard  
Manassas, Virginia 20110-2209  
United States of America

Date of deposit

11 April 2001

Accession Number

Unassigned

**C. ADDITIONAL INDICATIONS** (leave blank if not applicable)

This information is continued on an additional sheet ☐

**D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE** (if the indications are not for all designated States)

Europe

In respect of those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28(4) EPC).

Continued on additional sheets

**E. SEPARATE FURNISHING OF INDICATIONS** (leave blank if not applicable)

The indications listed below will be submitted to the international Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")

For receiving Office use only		For International Bureau use only	
<input checked="" type="checkbox"/> This sheet was received with the international application <b>Enora Rivera</b> <b>PCT Operations - IAPD Team 1</b> Authorized officer (703) 305-3678 (703) 305-3230 (703) 305-3230		<input type="checkbox"/> This sheet was received by the International Bureau on:  Authorized officer	

**ATCC Deposit No.: Unassigned**

## **CANADA**

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

## **NORWAY**

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

## **AUSTRALIA**

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

## **FINLAND**

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

## **UNITED KINGDOM**

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

**ATCC Deposit No.: Unassigned**

## **DENMARK**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

## **SWEDEN**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

## **NETHERLANDS**

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

**What is claimed:**

1. An albumin fusion protein comprising a Therapeutic protein:X and albumin comprising the amino acid sequence of SEQ ID NO:18.
2. An albumin fusion protein comprising a Therapeutic protein:X and a fragment or a variant of the amino acid sequence of SEQ ID NO:18, wherein said fragment or variant has albumin activity.
3. The albumin fusion protein of claim 2, wherein said albumin activity is the ability to prolong the shelf life of the Therapeutic protein:X compared to the shelf-life of the Therapeutic protein:X in an unfused state.
4. The albumin fusion protein of claim 2, wherein the fragment or variant comprises the amino acid sequence of amino acids 1-387 of SEQ ID NO:18.
5. The albumin fusion protein of any one of claims 1-4, wherein said Therapeutic protein:X comprises interferon-alpha.
6. An albumin fusion protein comprising a fragment or variant of a Therapeutic protein:X, and albumin comprising the amino acid sequence of SEQ ID NO:18, wherein said fragment or variant has a biological activity of the Therapeutic protein:X.
7. The albumin fusion protein of claim 6, wherein said Therapeutic protein:X comprises interferon-alpha, and wherein said fragment or variant has antiviral activity or inhibits cell proliferation.
8. The albumin fusion protein of any one of claims 1-4 or 6, wherein said Therapeutic protein:X, or fragment or variant thereof, comprises a protein selected from the

group consisting of:

- (a) serum cholinesterase;
- (b) alpha-1 antitrypsin;
- (c) aprotinin;
- (d) coagulation complex;
- (e) von Willebrand factor;
- (f) fibrinogen;
- (g) factor VII;
- (h) factor VIIA activated factor;
- (i) factor VIII;
- (j) factor IX;
- (k) factor X;
- (l) factor XIII;
- (m) c1 inactivator;
- (n) antithrombin III;
- (o) thrombin;
- (p) prothrombin;
- (q) apo-lipoprotein;
- (r) c-reactive protein;
- (s) protein C; and
- (t) immunoglobulin.

9. The albumin fusion protein of any one of claims 1-8, wherein the Therapeutic protein:X, or fragment or variant thereof, is fused to the N-terminus of albumin, or the N-terminus of the fragment or variant of albumin.

10. The albumin fusion protein of any one of claims 1-8, wherein the Therapeutic protein:X, or fragment or variant thereof, is fused to the C-terminus of albumin, or the C-terminus of the fragment or variant of albumin.



11. The albumin fusion protein of any one of claims 1-8, wherein the Therapeutic protein:X, or fragment or variant thereof, is fused to the N- terminus and C-terminus of albumin, or the N-terminus and the C-terminus of the fragment or variant of albumin.

12. The albumin fusion protein of any one of claims 1-8, which comprises a first Therapeutic protein:X, or fragment or variant thereof, and a second Therapeutic protein:X, or fragment or variant thereof, wherein said first Therapeutic protein:X, or fragment or variant thereof, is different from said second Therapeutic protein:X, or fragment or variant thereof.

13. The albumin fusion protein of any one of claims 1-11, wherein the Therapeutic protein:X, or fragment or variant thereof, is separated from the albumin or the fragment or variant of albumin by a linker.

14. The albumin fusion protein of any one of claims 1-11, wherein the albumin fusion protein has the following formula:

$R1-L-R2$ ;  $R2-L-R1$ ; or  $R1-L-R2-L-R1$ ,

wherein R1 is Therapeutic protein:X, or fragment or variant thereof, L is a peptide linker, and R2 is albumin comprising the amino acid sequence of SEQ ID NO:18 or fragment or variant of albumin.

15. The albumin fusion protein of any one of claims 1-14, wherein the shelf-life of the albumin fusion protein is greater than the shelf-life of the Therapeutic protein:X in an unfused state.

16. The albumin fusion protein of any one of claims 1-14, wherein the in vitro biological activity of the Therapeutic protein:X, or fragment or variant thereof, fused to

albumin, or fragment or variant thereof, is greater than the in vitro biological activity of the Therapeutic protein:X, or fragment or variant thereof, in an unfused state.

17. The albumin fusion protein of any one of claims 1-14, wherein the in vivo biological activity of the Therapeutic protein:X, or fragment or variant thereof, fused to albumin, or fragment or variant thereof, is greater than the in vivo biological activity of the Therapeutic protein:X, or fragment or variant thereof, in an unfused state.

18. An albumin fusion protein comprising a peptide inserted into an albumin comprising the amino acid sequence of SEQ ID NO:18 or fragment or variant thereof.

19. An albumin fusion protein comprising a peptide inserted into an albumin comprising an amino acid sequence selected from the group consisting of:

- (a) amino acids 54 to 61 of SEQ ID NO:18;
- (b) amino acids 76 to 89 of SEQ ID NO:18;
- (c) amino acids 92 to 100 of SEQ ID NO:18;
- (d) amino acids 170 to 176 of SEQ ID NO:18;
- (e) amino acids 247 to 252 of SEQ ID NO:18;
- (f) amino acids 266 to 277 of SEQ ID NO:18;
- (g) amino acids 280 to 288 of SEQ ID NO:18;
- (h) amino acids 362 to 368 of SEQ ID NO:18;
- (i) amino acids 439 to 447 of SEQ ID NO:18;
- (j) amino acids 462 to 475 of SEQ ID NO:18;
- (k) amino acids 478 to 486 of SEQ ID NO:18; and
- (l) amino acids 560 to 566 of SEQ ID NO:18.

20. The albumin fusion protein of claims 18 or 19, wherein said albumin fusion protein comprises a portion of albumin sufficient to prolong the shelf-life of the peptide as compared to the shelf-life of the peptide in an unfused state.

21. The albumin fusion protein of claims 18 or 19, wherein said albumin fusion protein comprises a portion of albumin sufficient to prolong the in vitro biological activity of the peptide fused to albumin as compared to the in vitro biological activity of the peptide in an unfused state.

22. The albumin fusion protein of claims 18 or 19 wherein said albumin fusion protein comprises a portion of albumin sufficient to prolong the in vivo biological activity of the peptide fused to albumin compared to the in vivo biological activity of the peptide in an unfused state.

23. An albumin fusion protein comprising a single chain antibody or portion thereof and albumin comprising the amino acid sequence of SEQ ID NO:18 or fragment or variant thereof.

24. The albumin fusion protein of any one of claims 1-23, which is non-glycosylated.

25. The albumin fusion protein of any one of claims 1-23, which is expressed in yeast.

26. The albumin fusion protein of claim 25, wherein the yeast is glycosylation deficient.

27. The albumin fusion protein of claim 25 wherein the yeast is glycosylation and protease deficient.

28. The albumin fusion protein of any one of claims 1-23, which is expressed by a mammalian cell.

29. The albumin fusion protein of any one of claims 1-23, wherein the albumin fusion protein is expressed by a mammalian cell in culture.

30. The albumin fusion protein of any one of claims 1-23, wherein the albumin fusion protein further comprises a secretion leader sequence.

31. A composition comprising the albumin fusion protein of any one of claims 1-30 and a pharmaceutically acceptable carrier.

32. A kit comprising the composition of claim 31.

33. A method of treating a disease or disorder in a patient, comprising the step of administering the albumin fusion protein of any one of claims 1-30.

34. The method of claim 33, wherein the disease or disorder comprises indication:Y.

35. The method of claim 34, wherein the Therapeutic protein:X comprises interferon-alpha, or fragment or variant thereof, and the disease or disorder is selected from the group consisting of: Hairy cell leukemia; Kaposi's sarcoma; genital warts; anal warts; chronic hepatitis B; chronic non-A, non-B hepatitis; hepatitis C; hepatitis D; chronic myelogenous leukemia; renal cell carcinoma; bladder carcinoma; ovarian carcinoma; cervical carcinoma; skin cancer; recurrent respirator papillomatosis; non-Hodgkin's lymphoma; cutaneous T-cell lymphoma; melanoma; multiple myeloma; AIDS; multiple sclerosis; and glioblastoma.

36. A method of treating a patient with a disease or disorder that is modulated by Therapeutic protein:X, comprising the step of administering an effective amount of the

albumin fusion protein of any one of claims 1-30.

37. The method of claim 36, wherein the disease or disorder is indication:Y.

38. The method of claim 37, wherein the Therapeutic protein:X is interferon-alpha, or fragment or variant thereof, and the disease or disorder is selected from the group consisting of: Hairy cell leukemia; Kaposi's sarcoma; genital warts; anal warts; chronic hepatitis B; chronic non-A, non-B hepatitis; hepatitis C; hepatitis D; chronic myelogenous leukemia; renal cell carcinoma; bladder carcinoma; ovarian carcinoma; cervical carcinoma; skin cancer; recurrent respirator papillomatosis; non-Hodgkin's lymphoma; cutaneous T-cell lymphoma; melanoma; multiple myeloma; AIDS; multiple sclerosis; and glioblastoma.


39. A method of extending the shelf life of Therapeutic protein:X comprising the step of fusing the Therapeutic protein:X, or fragment or variant thereof, to albumin or a fragment or variant of albumin sufficient to extend the shelf-life of the Therapeutic protein:X, or fragment or variant thereof, compared to the shelf-life of the Therapeutic protein:X, or fragment or variant thereof, in an unfused state.

40. A nucleic acid molecule comprising a polynucleotide sequence encoding the albumin fusion protein of any one of claims 1-30.


41. A vector comprising the nucleic acid molecule of claim 40.

42. A host cell comprising the nucleic acid molecule of claim 40.



1/20

**Figure 1**

2/20

**Figure 2**

3/20

**Figure 3A****Figure 3B**



4/20





Figure 4

5/20

**Figure 5**

6/20




Figure 6

7/20





Figure 7

8/20

**FIG. 8**

9/20

**Localisation of 'Loops' based on the HA Crvstal Structure**  
**which could be used for Mutation/Insertion**

```

1    DAHKSEVAHR FKDLGEENFK ALVLIAFAQY LQQCPFEDHV KLVNEVTEFA
      HHHHH HHH      HHH HHHHHHHHHHH      HHHHH HHHHHHHHHHH

      I                      II                      III
51   KTCVADESAE NCDKSLHTLF GDKLCTVATL RETYGEMADC CAKOEPERNE
      HHHHH      HHHHH HHHHH      HHHH H      HHHH

101  CFLQHKDDNP NLPRLVRPEV DVMCTAFHDN EETFLKKYLY EIARRHPYFY
      HHHH      H      HHHHHHHHH      HHHHHHHHH HHHHH

      IV
151  APELLFFAKR YKAAFTCCO AADKAACLLP KLDEL RDEGK ASSAKQRLKC
      HHHHHHHHHHH HHHHHHHHH      HHHHH HHHHHHHHHHH HHHHHHHHHHH

      V
201  ASLQKFGERA FKAWAVARLS QRFPAEFAE VSKLVTDLTK VHTECCHGDL
      HHHHH      HH HHHHHHHHHHH HH      HHH HHHHHHHHHHH HHHHHH      HH

      VI                      VII
251  LECADDRADL AKYICENODS ISSKLKECCE KPLLEKSHCI AEVENDEMPA
      HHHHHHHHHHH HHHHH      HHHHH      HHHHHHH H

301  DLPSLAADFV ESKDVCKNYA EAKDVFLGMF LYEYARRHPD YSVVLLRLA
      HHHH      HHHHHH      HHHHHHH HHHHHH      HHHHHHHHH

      VIII
351  KTYETTLEKC CAAADPHECY AKVFDEFKPL VEEPQNLIKQ NCELFEQLGE
      HHHHHHHHHHH      HH      H      HHHHH HHHHHHHHHHH HHHHHHH

      IX
401  YKFQNALLVR YTKKVPQVST PTLVEVSRNL GKVGSKCCKH PEAKRMPCAE
      HHHHHHHHHHH HHHH      H HHHHHHHHHHH      HHH      HHHHHHHHH

      X                      XI
451  DYLSVVLNQL CVLHEKTPVS DRVTKCCTES LVNRRPPCFSA LEVDETYVPK
      HHHHHHHHHHH HHHHH      HHHHHHHHH      HHHHHHHHH

501  EFNAETFTFH ADICTLSEKE RQIKKQTALV ELVKHKPKAT KEQLKAVMDD
      HHH      HHH HHHHHMEHHH HHH      HHHHHHHHH

      XII
551  FAAFVEKCK ADDKETCFAE EGKKLVAASQ AALGL
      HHHHHHHHH      HHHH HHHHHHHHHHH HH

```

**Loop**

I Val54-Asn61  
 II Thr76-Asp89  
 III Ala92-Glu100  
 IV Gln170-Ala176  
 V His247-Glu252  
 VI Glu266-Glu277

**Loop**

VII Glu280-His288  
 VIII Ala362-Glu368  
 IX Lys439-Pro447  
 X Val462-Lys475  
 XI Thr478-Pro486  
 XII Lys560-Thr566

**Figure 9**

10/20

**Examples of Modifications to Loop IV****a. Randomisation of Loop IV.**

IV

```

151  APELLFFAKR YKAAFTECCQ AADKAACLLP KLDEL RDEGK ASSAKQRLKC
      HHHHHHHHHH HHHHHHHHHH          HHHHH HHHHHHHHHHH HHHHHHHHHHH

```

IV

```

151  APELLFFAKR YKAAFTECCX XXXXXXCLLP KLDEL RDEGK ASSAKQRLKC
      HHHHHHHHHH HHHHHHHHHH          HHHHH HHHHHHHHHHH HHHHHHHHHHH

```

**X** represents the mutation of the natural amino acid to any other amino acid. One, more or all of the amino acids can be changed in this manner. This figure indicates all the residues have been changed.

**b. Insertion (or replacement) of Randomised sequence into Loop IV.**

(X)<sub>n</sub>  
↓  
IV

```

151  APELLFFAKR YKAAFTECCQ AADKAACLLP KLDEL RDEGK ASSAKQRLKC
      HHHHHHHHHH HHHHHHHHHH          HHHHH HHHHHHHHHHH HHHHHHHHHHH

```

The insertion can be at any point on the loop and the length a length where n would typically be 6, 8, 12, 20 or 25.

**Figure 10**





FIG. 11A






**FIG. 11B**

FROM FIG. 11B


FROM FIG. 11B

13/20




**FIG. 11C**

14/20



**FIG. 12:**  
LOOP IV GLU170-A176

15/20



**FIG. 13**  
TERTIARY STRUCTURE OF HA

16/20




FIG. 14

17/20

```

1  GAT GCA CAC AAG AGT GAG GTT GCT CAT CGG TTT AAA GAT TTG GGA GAA AAT TTC AAA 60
1  D  A  H  K  S  E  V  A  H  R  F  K  D  L  G  E  E  N  F  K  20

61  GCC TTG GTG TTG ATT GCC TTT GCT CAG TAT CTT CAG CAG TGT CCA TTT GAA GAT CAT GTA 120
21  A  L  V  L  I  A  F  A  Q  Y  L  Q  Q  C  P  F  E  D  H  V  40

121 AAA TTA GTG AAT GAA GTA ACT GAA TTT GCA AAA ACA TGT GTT GCT GAT GAG TCA GCT GAA 180
41  K  L  V  N  E  V  T  E  F  A  K  T  C  V  A  D  E  S  A  E  60

181 AAT TGT GAC AAA TCA CTT CAT ACC CTT TTT GGA GAC AAA TTA TGC ACA GTT GCA ACT CTT 240
61  N  C  D  K  S  L  H  T  L  F  G  D  K  L  C  T  V  A  T  L  80

241 CGT GAA ACC TAT GGT GAA ATG GCT GAC TGC TGT GCA AAA CAA GAA CCT GAG AGA AAT GAA 300
81  R  E  T  Y  G  E  M  A  D  C  C  A  K  Q  E  P  E  R  N  E  100

301 TGC TTC TTG CAA CAC AAA GAT GAC AAC CCA AAC CTC CCC CGA TTG GTG AGA CCA GAG GTT 360
101 C  F  L  Q  H  K  D  D  N  P  N  L  P  R  L  V  R  P  E  V  120

361 GAT GTG ATG TGC ACT GCT TTT CAT GAC AAT GAA GAG ACA TTT TTG AAA AAA TAC TTA TAT 420
121 D  V  M  C  T  A  F  H  D  N  E  E  T  F  L  K  K  Y  L  Y  140

421 GAA ATT GCC AGA AGA CAT CCT TAC TTT TAT GCC CCG GAA CTC CTT TTC TTT GCT AAA AGG 480
141 E  I  A  R  R  H  P  Y  F  Y  A  P  E  L  L  F  F  A  K  R  160

```

Figure 15A

18/20

```

481 TAT AAA GCT GCT TTT ACA GAA TGT TGC CAA GCT GCT GAT AAA GCT GCC TGC CTG TTG CCA 540
161 Y K A A F T E C C Q A A D K A A C L L P 180

541 AAG CTC GAT GAA CTT CGG GAT GAA GGG AAG GCT TCG TCT GCC AAA CAG AGA CTC AAA TGT 600
181 K L D E L R D E G K A S S A K Q R L K C 200

601 GCC AGT CTC CAA AAA TTT GGA GAA AGA GCT TTC AAA GCA TGG GCA GTG GCT CGC CTG AGC 660
201 A S L Q K K F G E R A F K A W A V A R L S 220

661 CAG AGA TTT CCC AAA GCT GAG TTT GCA GAA GTT TCC AAG TTA GTG ACA GAT CTT ACC AAA 720
221 Q R F P K A E F A E V S K L V T D L T K 240

721 GTC CAC ACG GAA TGC TGC CAT GGA GAT CTG CTT GAA TGT GCT GAT GAC AGG GCG GAC CTT 780
241 V H T E C C H G D L L E C A D D R A D L 260

781 GCC AAG TAT ATC TGT GAA AAT CAG GAT TCG ATC TCC AGT AAA CTG AAG GAA TGC TGT GAA 840
261 A K Y I C E N Q D S I S S K L K E C E 280

841 AAA CCT CTG TTG GAA AAA TCC CAC TGC ATT GCC GAA GTG GAA AAT GAT GAG ATG CCT GCT 900
281 K P L L E K S H C I A E V E N D E M P A 300

901 GAC TTG CCT TCA TTA GCT GCT GAT TTT GTT GAA AGT AAG GAT GTT TGC AAA AAC TAT GCT 960
301 D L P S L A A A D F V E S K D V C K N Y A 320

```

Figure 15B

19/20

```

961 GAG GCA AAG GAT GTC TTC CTG GGC ATG TTT TTG TAT GAA TAT GCA AGA AGG CAT CCT GAT 1020
321 E A K D V F L G M F L Y E Y A R R H P D 340

1021 TAC TCT GTC GTG CTG CTG GAT CAT GAA TGC TAT GCC AAG ACA TAT GAA ACC ACT CTA GAG AAG TGC 1080
341 Y S V V L L L L R L A K T Y E T T L E K C 360

1081 TGT GCC GCT GCA GAT CCT CAT GAA TGC TAT GCC AAA GTG TTC GAT GAA TTT AAA CCT CTT 1140
361 C A A A D P H E C Y A K V F D E F K P L 380

1141 GTG GAA GAG CCT CAG AAT TTA ATC AAA CAA AAC TGT GAG CTT TTT GAG CAG CTT GGA GAG 1200
381 V E E P Q N L I K Q N C E L F E Q L G E 400

1201 TAC AAA TTC CAG AAT GCG CTA TTA GTT CGT TAC ACC AAG AAA GTA CCC CAA GTG TCA ACT 1260
401 Y K F Q N A L L V R Y T K K V P Q V S T 420

1261 CCA ACT CTT GTA GAG GTC TCA AGA AAC CTA GGA AAA GTG GGC AGC AAA TGT TGT AAA CAT 1320
421 P T L V E V S S R N L G K V G S K C C K H 440

1321 CCT GAA GCA AAA AGA ATG CCC TGT GCA GAA GAC TAT CTA TCC GTG GTC CTG AAC CAG TTA 1380
441 P E A K R M P C A E D Y L S V V L N Q L 460

1381 TGT GTG TTG CAT GAG AAA ACG CCA GTA AGT GAC AGA GTC ACA AAA TGC TGC ACA GAG TCC 1440
461 C V L H E K T P V S D R V T K C C T E S 480

```

Figure 15C



20/20

```

1441 TTG GTG AAC AGG CGA CCA TGC TTT TCA GCT CTG GAA GTC GAT GAA ACA TAC GTT CCC AAA 1500
481 L V N R R R P C F S A L E V D E T Y V P K 500

1501 GAG TTT AAT GCT GAA ACA TTC ACC TTC CAT GCA GAT ATA TGC ACA CTT TCT GAG AAG GAG 1560
501 E F N A E T F T F H A D I C T L S E K E 520

1561 AGA CAA ATC AAG AAA CAA ACT GCA CTT GTT GAG CTT GTG AAA CAC AAG CCC AAG GCA ACA 1620
521 R Q I K K Q T A L V E L V K H K P K A T 540

1621 AAA GAG CAA CTG AAA GCT GTT ATG GAT GAT TTC GCA GCT TTT GTA GAG AAG TGC TGC AAG 1680
541 K E Q L K A V M D D F A A F V E K C C K 560

1681 GCT GAC GAT AAG GAG ACC TGC TTT GCC GAG GAG GGT AAA AAA CTT GTT GCT GCA AGT CAA 1740
561 A D D K E T C F A E E G K K L V A A S Q 580

1741 GCT GCC TTA GGC TTA TAA CAT CTA CAT TTA AAA GCA TCT CAG 1782
581 A A L G L * 585

```

Figure 15D

## SEQUENCE LISTING

<110> Delta Biotechnology Limited  
Principia Pharmaceutical Corporation

<120> Albumin Fusion Proteins

<130> PF542PCT

<140> Unassigned

<141> 2001-04-12

<150> 60/229,358

<151> 2000-04-12

<150> 60/256,931

<151> 2000-12-21

<150> 60/199,384

<151> 2000-04-25

<160> 37

<170> PatentIn Ver. 2.1

<210> 1

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<221> primer\_bind

<223> primer useful to clone human growth hormone cDNA

<400> 1

ccaagaatt cccttatcca ggc

23

<210> 2

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<221> primer\_bind

<223> primer useful to clone human growth hormone cDNA

<400> 2

gggaagctta gaagccacag gatccctcca cag

33

<210> 3

<211> 16

<212> DNA

<213> Artificial Sequence

<220>

<221> misc\_structure  
<223> synthetic oligonucleotide used to join DNA fragments  
with non-cohesive ends.

<400> 3  
gataaagatt cccaac 16

<210> 4  
<211> 17  
<212> DNA  
<213> Artificial Sequence

<220>  
<221> misc\_structure  
<223> synthetic oligonucleotide used to join DNA fragments  
with non-cohesive ends.

<400> 4  
aattgttggg aatcttt 17

<210> 5  
<211> 17  
<212> DNA  
<213> Artificial Sequence

<220>  
<221> misc\_structure  
<223> synthetic oligonucleotide used to join DNA fragments  
with non-cohesive ends.

<400> 5  
ttaggcttat tccaac 17

<210> 6  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>  
<221> misc\_structure  
<223> synthetic oligonucleotide used to join DNA fragments  
with non-cohesive ends.

<400> 6  
aattgttggg aataagcc 18

<210> 7  
<211> 24  
<212> PRT  
<213> Artificial Sequence

<220>  
<221> SITE  
<222> 1)..(19)  
<223> invertase leader sequence

<220>  
 <221> SITE  
 <222> 20)..(24)  
 <223> first 5 amino acids of mature human serum albumin

<400> 7  
 Met Leu Leu Gln Ala Phe Leu Phe Leu Leu Ala Gly Phe Ala Ala Lys  
           1                  5                          10                          15  
 Ile Ser Ala Asp Ala His Lys Ser  
                           20

<210> 8  
 <211> 21  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> misc\_structure  
 <223> synthetic oligonucleotide used to join DNA  
 fragments with non-cohesive ends.

<400> 8  
 gagatgcaca cctgagtgag g 21

<210> 9  
 <211> 27  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> misc\_structure  
 <223> synthetic oligonucleotide used to join DNA  
 fragments with non-cohesive ends.

<400> 9  
 gatcctgtgg cttcgatgca cacaaga 27

<210> 10  
 <211> 24  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> misc\_structure  
 <223> synthetic oligonucleotide used to join DNA  
 fragments with non-cohesive ends.

<400> 10  
 ctcttgtgtg catcgaagcc acag 24

<210> 11  
 <211> 30  
 <212> DNA  
 <213> Artificial Sequence

<220>  
<221> misc\_structure  
<223> synthetic oligonucleotide used to join DNA fragments with non-cohesive ends.

<400> 11  
tgtggaagag cctcagaatt tattcccaac 30

<210> 12  
<211> 31  
<212> DNA  
<213> Artificial Sequence

<220>  
<221> misc\_structure  
<223> synthetic oligonucleotide used to join DNA fragments with non-cohesive ends.

<400> 12  
aattggtggg aataaattct gaggctcttc c 31

<210> 13  
<211> 47  
<212> DNA  
<213> Artificial Sequence

<220>  
<221> misc\_structure  
<223> synthetic oligonucleotide used to join DNA fragments with non-cohesive ends.

<400> 13  
ttaggcttag gtggcggtgg atccggcggt ggtggatctt tcccaac 47

<210> 14  
<211> 48  
<212> DNA  
<213> Artificial Sequence

<220>  
<221> misc\_structure  
<223> synthetic oligonucleotide used to join DNA fragments with non-cohesive ends.

<400> 14  
aattggtggg aaagatccac caccgccgga tccaccgcca cctaagcc 48

<210> 15  
<211> 62  
<212> DNA  
<213> Artificial Sequence

<220>  
<221> misc\_structure  
<223> synthetic oligonucleotide used to join DNA fragments with non-cohesive ends.

&lt;400&gt; 15

ttaggcttag gcggtggtgg atctggtggc ggcggatctg gtggcggtgg atccttccca 60  
ac 62

&lt;210&gt; 16

&lt;211&gt; 63

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;221&gt; misc\_structure

<223> synthetic oligonucleotide used to join DNA  
fragments with non-cohesive ends.

&lt;400&gt; 16

aattgttggg aaggatccac cgccaccaga tccgcccga ccagatccac caccgcctaa 60  
gcc 63

&lt;210&gt; 17

&lt;211&gt; 1782

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(1755)

&lt;400&gt; 17

gat gca cac aag agt gag gtt gct cat cgg ttt aaa gat ttg gga gaa 48  
Asp Ala His Lys Ser Glu Val Ala His Arg Phe Lys Asp Leu Gly Glu  
1 5 10 15

gaa aat ttc aaa gcc ttg gtg ttg att gcc ttt gct cag tat ctt cag 96  
Glu Asn Phe Lys Ala Leu Val Leu Ile Ala Phe Ala Gln Tyr Leu Gln  
20 25 30

cag tgt cca ttt gaa gat cat gta aaa tta gtg aat gaa gta act gaa 144  
Gln Cys Pro Phe Glu Asp His Val Lys Leu Val Asn Glu Val Thr Glu  
35 40 45

ttt gca aaa aca tgt gtt gct gat gag tca gct gaa aat tgt gac aaa 192  
Phe Ala Lys Thr Cys Val Ala Asp Glu Ser Ala Glu Asn Cys Asp Lys  
50 55 60

tca ctt cat acc ctt ttt gga gac aaa tta tgc aca gtt gca act ctt 240  
Ser Leu His Thr Leu Phe Gly Asp Lys Leu Cys Thr Val Ala Thr Leu  
65 70 75 80

cgt gaa acc tat ggt gaa atg gct gac tgc tgt gca aaa caa gaa cct 288  
Arg Glu Thr Tyr Gly Glu Met Ala Asp Cys Cys Ala Lys Gln Glu Pro  
85 90 95

gag aga aat gaa tgc ttc ttg caa cac aaa gat gac aac cca aac ctc 336  
Glu Arg Asn Glu Cys Phe Leu Gln His Lys Asp Asp Asn Pro Asn Leu  
100 105 110

ccc cga ttg gtg aga cca gag gtt gat gtg atg tgc act gct ttt cat	384
Pro Arg Leu Val Arg Pro Glu Val Asp Val Met Cys Thr Ala Phe His	
115 120 125	
 gac aat gaa gag aca ttt ttg aaa aaa tac tta tat gaa att gcc aga	432
Asp Asn Glu Glu Thr Phe Leu Lys Lys Tyr Leu Tyr Glu Ile Ala Arg	
130 135 140	
 aga cat cct tac ttt tat gcc ccg gaa ctc ctt ttc ttt gct aaa agg	480
Arg His Pro Tyr Phe Tyr Ala Pro Glu Leu Leu Phe Phe Ala Lys Arg	
145 150 155 160	
 tat aaa gct gct ttt aca gaa tgt tgc caa gct gct gat aaa gct gcc	528
Tyr Lys Ala Ala Phe Thr Glu Cys Cys Gln Ala Ala Asp Lys Ala Ala	
165 170 175	
 tgc ctg ttg cca aag ctc gat gaa ctt cgg gat gaa ggg aag gct tcg	576
Cys Leu Leu Pro Lys Leu Asp Glu Leu Arg Asp Glu Gly Lys Ala Ser	
180 185 190	
 tct gcc aaa cag aga ctc aaa tgt gcc agt ctc caa aaa ttt gga gaa	624
Ser Ala Lys Gln Arg Leu Lys Cys Ala Ser Leu Gln Lys Phe Gly Glu	
195 200 205	
 aga gct ttc aaa gca tgg gca gtg gct cgc ctg agc cag aga ttt ccc	672
Arg Ala Phe Lys Ala Trp Ala Val Ala Arg Leu Ser Gln Arg Phe Pro	
210 215 220	
 aaa gct gag ttt gca gaa gtt tcc aag tta gtg aca gat ctt acc aaa	720
Lys Ala Glu Phe Ala Glu Val Ser Lys Leu Val Thr Asp Leu Thr Lys	
225 230 235 240	
 gtc cac acg gaa tgc tgc cat gga gat ctg ctt gaa tgt gct gat gac	768
Val His Thr Glu Cys Cys His Gly Asp Leu Leu Glu Cys Ala Asp Asp	
245 250 255	
 agg gcg gac ctt gcc aag tat atc tgt gaa aat cag gat tcg atc tcc	816
Arg Ala Asp Leu Ala Lys Tyr Ile Cys Glu Asn Gln Asp Ser Ile Ser	
260 265 270	
 agt aaa ctg aag gaa tgc tgt gaa aaa cct ctg ttg gaa aaa tcc cac	864
Ser Lys Leu Lys Glu Cys Cys Glu Lys Pro Leu Leu Glu Lys Ser His	
275 280 285	
 tgc att gcc gaa gtg gaa aat gat gag atg cct gct gac ttg cct tca	912
Cys Ile Ala Glu Val Glu Asn Asp Glu Met Pro Ala Asp Leu Pro Ser	
290 295 300	
 tta gct gct gat ttt gtt gaa agt aag gat gtt tgc aaa aac tat gct	960
Leu Ala Ala Asp Phe Val Glu Ser Lys Asp Val Cys Lys Asn Tyr Ala	
305 310 315 320	
 gag gca aag gat gtc ttc ctg ggc atg ttt ttg tat gaa tat gca aga	1008
Glu Ala Lys Asp Val Phe Leu Gly Met Phe Leu Tyr Glu Tyr Ala Arg	
325 330 335	
 agg cat cct gat tac tct gtc gtg ctg ctg ctg aga ctt gcc aag aca	1056
Arg His Pro Asp Tyr Ser Val Val Leu Leu Leu Arg Leu Ala Lys Thr	
340 345 350	

tat gaa acc act cta gag aag tgc tgt gcc gct gca gat cct cat gaa	1104
Tyr Glu Thr Thr Leu Glu Lys Cys Cys Ala Ala Ala Asp Pro His Glu	
355 360 365	
tgc tat gcc aaa gtg ttc gat gaa ttt aaa cct ctt gtg gaa gag cct	1152
Cys Tyr Ala Lys Val Phe Asp Glu Phe Lys Pro Leu Val Glu Glu Pro	
370 375 380	
cag aat tta atc aaa caa aac tgt gag ctt ttt gag cag ctt gga gag	1200
Gln Asn Leu Ile Lys Gln Asn Cys Glu Leu Phe Glu Gln Leu Gly Glu	
385 390 395 400	
tac aaa ttc cag aat gcg cta tta gtt cgt tac acc aag aaa gta ccc	1248
Tyr Lys Phe Gln Asn Ala Leu Leu Val Arg Tyr Thr Lys Lys Val Pro	
405 410 415	
caa gtg tca act cca act ctt gta gag gtc tca aga aac cta gga aaa	1296
Gln Val Ser Thr Pro Thr Leu Val Glu Val Ser Arg Asn Leu Gly Lys	
420 425 430	
gtg ggc agc aaa tgt tgt aaa cat cct gaa gca aaa aga atg ccc tgt	1344
Val Gly Ser Lys Cys Cys Lys His Pro Glu Ala Lys Arg Met Pro Cys	
435 440 445	
gca gaa gac tat cta tcc gtg gtc ctg aac cag tta tgt gtg ttg cat	1392
Ala Glu Asp Tyr Leu Ser Val Val Leu Asn Gln Leu Cys Val Leu His	
450 455 460	
gag aaa acg cca gta agt gac aga gtc aca aaa tgc tgc aca gag tcc	1440
Glu Lys Thr Pro Val Ser Asp Arg Val Thr Lys Cys Cys Thr Glu Ser	
465 470 475 480	
ttg gtg aac agg cga cca tgc ttt tca gct ctg gaa gtc gat gaa aca	1488
Leu Val Asn Arg Arg Pro Cys Phe Ser Ala Leu Glu Val Asp Glu Thr	
485 490 495	
tac gtt ccc aaa gag ttt aat gct gaa aca ttc acc ttc cat gca gat	1536
Tyr Val Pro Lys Glu Phe Asn Ala Glu Thr Phe Thr Phe His Ala Asp	
500 505 510	
ata tgc aca ctt tct gag aag gag aga caa atc aag aaa caa act gca	1584
Ile Cys Thr Leu Ser Glu Lys Glu Arg Gln Ile Lys Lys Gln Thr Ala	
515 520 525	
ctt gtt gag ctt gtg aaa cac aag ccc aag gca aca aaa gag caa ctg	1632
Leu Val Glu Leu Val Lys His Lys Pro Lys Ala Thr Lys Glu Gln Leu	
530 535 540	
aaa gct gtt atg gat gat ttc gca gct ttt gta gag aag tgc tgc aag	1680
Lys Ala Val Met Asp Asp Phe Ala Ala Phe Val Glu Lys Cys Cys Lys	
545 550 555 560	
gct gac gat aag gag acc tgc ttt gcc gag gag ggt aaa aaa ctt gtt	1728
Ala Asp Asp Lys Glu Thr Cys Phe Ala Glu Glu Gly Lys Lys Leu Val	
565 570 575	
gct gca agt caa gct gcc tta ggc tta taacatctac atttaaaagc atctcag	1782
Ala Ala Ser Gln Ala Ala Leu Gly Leu	
580 585	



&lt;210&gt; 18

&lt;211&gt; 585

&lt;212&gt; PRT

&lt;213&gt; Homo Sapiens

&lt;400&gt; 18

Asp Ala His Lys Ser Glu Val Ala His Arg Phe Lys Asp Leu Gly Glu  
 1 5 10 15

Glu Asn Phe Lys Ala Leu Val Leu Ile Ala Phe Ala Gln Tyr Leu Gln  
 20 25 30

Gln Cys Pro Phe Glu Asp His Val Lys Leu Val Asn Glu Val Thr Glu  
 35 40 45

Phe Ala Lys Thr Cys Val Ala Asp Glu Ser Ala Glu Asn Cys Asp Lys  
 50 55 60

Ser Leu His Thr Leu Phe Gly Asp Lys Leu Cys Thr Val Ala Thr Leu  
 65 70 75 80

Arg Glu Thr Tyr Gly Glu Met Ala Asp Cys Cys Ala Lys Gln Glu Pro  
 85 90 95

Glu Arg Asn Glu Cys Phe Leu Gln His Lys Asp Asp Asn Pro Asn Leu  
 100 105 110

Pro Arg Leu Val Arg Pro Glu Val Asp Val Met Cys Thr Ala Phe His  
 115 120 125

Asp Asn Glu Glu Thr Phe Leu Lys Lys Tyr Leu Tyr Glu Ile Ala Arg  
 130 135 140

Arg His Pro Tyr Phe Tyr Ala Pro Glu Leu Leu Phe Phe Ala Lys Arg  
 145 150 155 160

Tyr Lys Ala Ala Phe Thr Glu Cys Cys Gln Ala Ala Asp Lys Ala Ala  
 165 170 175

Cys Leu Leu Pro Lys Leu Asp Glu Leu Arg Asp Glu Gly Lys Ala Ser  
 180 185 190

Ser Ala Lys Gln Arg Leu Lys Cys Ala Ser Leu Gln Lys Phe Gly Glu  
 195 200 205

Arg Ala Phe Lys Ala Trp Ala Val Ala Arg Leu Ser Gln Arg Phe Pro  
 210 215 220

Lys Ala Glu Phe Ala Glu Val Ser Lys Leu Val Thr Asp Leu Thr Lys  
 225 230 235 240

Val His Thr Glu Cys Cys His Gly Asp Leu Leu Glu Cys Ala Asp Asp  
 245 250 255

Arg Ala Asp Leu Ala Lys Tyr Ile Cys Glu Asn Gln Asp Ser Ile Ser  
 260 265 270

Ser Lys Leu Lys Glu Cys Cys Glu Lys Pro Leu Leu Glu Lys Ser His  
 275 280 285

Cys Ile Ala Glu Val Glu Asn Asp Glu Met Pro Ala Asp Leu Pro Ser  
 290 295 300  
 Leu Ala Ala Asp Phe Val Glu Ser Lys Asp Val Cys Lys Asn Tyr Ala  
 305 310 315 320  
 Glu Ala Lys Asp Val Phe Leu Gly Met Phe Leu Tyr Glu Tyr Ala Arg  
 325 330 335  
 Arg His Pro Asp Tyr Ser Val Val Leu Leu Arg Leu Ala Lys Thr  
 340 345 350  
 Tyr Glu Thr Thr Leu Glu Lys Cys Cys Ala Ala Ala Asp Pro His Glu  
 355 360 365  
 Cys Tyr Ala Lys Val Phe Asp Glu Phe Lys Pro Leu Val Glu Glu Pro  
 370 375 380  
 Gln Asn Leu Ile Lys Gln Asn Cys Glu Leu Phe Glu Gln Leu Gly Glu  
 385 390 395 400  
 Tyr Lys Phe Gln Asn Ala Leu Leu Val Arg Tyr Thr Lys Lys Val Pro  
 405 410 415  
 Gln Val Ser Thr Pro Thr Leu Val Glu Val Ser Arg Asn Leu Gly Lys  
 420 425 430  
 Val Gly Ser Lys Cys Cys Lys His Pro Glu Ala Lys Arg Met Pro Cys  
 435 440 445  
 Ala Glu Asp Tyr Leu Ser Val Val Leu Asn Gln Leu Cys Val Leu His  
 450 455 460  
 Glu Lys Thr Pro Val Ser Asp Arg Val Thr Lys Cys Cys Thr Glu Ser  
 465 470 475 480  
 Leu Val Asn Arg Arg Pro Cys Phe Ser Ala Leu Glu Val Asp Glu Thr  
 485 490 495  
 Tyr Val Pro Lys Glu Phe Asn Ala Glu Thr Phe Thr Phe His Ala Asp  
 500 505 510  
 Ile Cys Thr Leu Ser Glu Lys Glu Arg Gln Ile Lys Lys Gln Thr Ala  
 515 520 525  
 Leu Val Glu Leu Val Lys His Lys Pro Lys Ala Thr Lys Glu Gln Leu  
 530 535 540  
 Lys Ala Val Met Asp Asp Phe Ala Ala Phe Val Glu Lys Cys Cys Lys  
 545 550 555 560  
 Ala Asp Asp Lys Glu Thr Cys Phe Ala Glu Glu Gly Lys Lys Leu Val  
 565 570 575  
 Ala Ala Ser Gln Ala Ala Leu Gly Leu  
 580 585

<210> 19  
<211> 57  
<212> DNA  
<213> Artificial Sequence

<220>  
<221> primer\_bind  
<223> primer used to generate XhoI and ClaI  
site in pPPC0006

<400> 19  
gcctcgagaa aagagatgca cacaagagtg aggttgctca tcgatttaaa gatttgg 57

<210> 20  
<211> 58  
<212> DNA  
<213> Artificial Sequence

<220>  
<221> primer\_bind  
<223> primer used in generation XhoI and ClaI  
site in pPPC0006

<400> 20  
aatcgatgag caacctcact cttgtgtgca tctcttttct cgaggctcct ggaataag 58

<210> 21  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<221> primer\_bind  
<223> primer used in generation XhoI and ClaI  
site in pPPC0006

<400> 21  
tacaaactta agagtccaat tagc 24

<210> 22  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<221> primer\_bind  
<223> primer used in generation XhoI and ClaI  
site in pPPC0006

<400> 22  
cacttctcta gagtggtttc atatgtctt 29

<210> 23  
<211> 60  
<212> DNA  
<213> Artificial Sequence

<220>

<221> Misc\_Structure

<223> Synthetic oligonucleotide used to alter restriction sites in pPPC0007

<400> 23

aagctgcctt aggcttataa taaggcgcgc cggccggccg tttaaactaa gcttaattct 60

<210> 24

<211> 60

<212> DNA

<213> Artificial Sequence

<220>

<221> Misc\_Structure

<223> Synthetic oligonucleotide used to alter restriction sites in pPPC0007

<400> 24

agaattaagc ttagtttaaa cggccggccg gcgcgcctta ttataagcct aaggcagctt 60

<210> 25

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<221> primer\_bind

<223> forward primer useful for generation of albumin fusion protein in which the albumin moiety is N-terminal of the Therapeutic Protein

<220>

<221> misc\_feature

<222> (18)

<223> n equals a,t,g, or c

<220>

<221> misc\_feature

<222> (19)

<223> n equals a,t,g, or c

<220>

<221> misc\_feature

<222> (20)

<223> n equals a,t,g, or c

<220>

<221> misc\_feature

<222> (21)

<223> n equals a,t,g, or c

<220>

<221> misc\_feature

<222> (22)

<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (23)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (24)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (25)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (26)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (27)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (28)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (29)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (30)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (31)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (32)  
<223> n equals a,t,g, or c

<400> 25  
aagctgcctt aggcttannn nnnnnnnnnn nn

32

<210> 26  
<211> 51  
<212> DNA  
<213> Artificial Sequence

<220>  
<221> primer\_bind

<223> reverse primer useful for generation of albumin fusion protein in which the albumin moiety is N-terminal of the Therapeutic Protein

<220>

<221> misc\_feature

<222> (37)

<223> n equals a,t,g, or c

<220>

<221> misc\_feature

<222> (38)

<223> n equals a,t,g, or c

<220>

<221> misc\_feature

<222> (39)

<223> n equals a,t,g, or c

<220>

<221> misc\_feature

<222> (40)

<223> n equals a,t,g, or c

<220>

<221> misc\_feature

<222> (41)

<223> n equals a,t,g, or c

<220>

<221> misc\_feature

<222> (42)

<223> n equals a,t,g, or c

<220>

<221> misc\_feature

<222> (43)

<223> n equals a,t,g, or c

<220>

<221> misc\_feature

<222> (44)

<223> n equals a,t,g, or c

<220>

<221> misc\_feature

<222> (45)

<223> n equals a,t,g, or c

<220>

<221> misc\_feature

<222> (46)

<223> n equals a,t,g, or c

<220>

<221> misc\_feature

<222> (47)

<223> n equals a,t,g, or c

<220>

<221> misc\_feature  
<222> (48)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (49)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (50)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (51)  
<223> n equals a,t,g, or c

<400> 26  
gcgcgcgttt aaacggccgg ccggcgcgcc ttattannnn nnnnnnnnnn n 51

<210> 27  
<211> 33  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> forward primer useful for generation of albumin fusion  
protein in which the albumin moiety is c-terminal of the  
Therapeutic Protein

<220>  
<221> misc\_feature  
<222> (19)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (20)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (21)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (22)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (23)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature

222> (24)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (25)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (26)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (27)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (28)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (29)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (30)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (31)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (32)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (33)  
<223> n equals a,t,g, or c

<400> 27  
aggagcgtcg acaaaagann nnnnnnnnnn nnn

33

<210> 28  
<211> 52  
<212> DNA  
<213> Artificial Sequence

<220>  
<221> primer\_bind  
<223> reverse primer useful for generation of albumin



fusion protein in which the albumin moiety is c-terminal of the Therapeutic Protein

<220>  
<221> misc\_feature  
<222> (38)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (39)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (40)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (41)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (42)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (43)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (44)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (45)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (46)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (47)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (48)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature

<222> (49)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (50)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (51)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (52)  
<223> n equals a,t,g, or c

<400> 28  
ctttaaatcg atgagcaacc tcactcttgt gtgcacnnnn nnnnnnnnnn nn 52

<210> 29  
<211> 24  
<212> PRT  
<213> Artificial Sequence

<220>  
<221> signal  
<223> signal peptide of natural human serum albumin protein

<400> 29  
Met Lys Trp Val Ser Phe Ile Ser Leu Leu Phe Leu Phe Ser Ser Ala  
1 5 10 15  
Tyr Ser Arg Ser Leu Asp Lys Arg  
20

<210> 30  
<211> 114  
<212> DNA  
<213> Artificial Sequence

<220>  
<221> primer\_bind  
<223> forward primer useful for generation of PC4:HSA  
albumin fusion VECTOR

<220>  
<221> misc\_feature  
<222> (5)..(10)  
<223> BamHI restriction site

<220>  
<221> misc\_feature  
<222> (11)..(16)  
<223> Hind III restriction site

<220>  
<221> misc\_feature

<222> (17)..(27)  
<223> Kozak sequence

<220>  
<221> misc\_feature  
<222> (25)..(97)  
<223> cds natural signal sequence of human serum albumin

<220>  
<221> misc\_feature  
<222> (75)..(81)  
<223> XhoI restriction site

<220>  
<221> misc\_feature  
<222> (98)..(114)  
<223> cds first six amino acids of human serum albumin

<400> 30  
tcaggggatcc aagcttcgcg caccatgaag tgggtaacct ttatttcctc tctttttctc 60  
ttagctcgg cttactcgag ggggtgtgtt cgtcgagatg cacacaagag tgag 114

<210> 31  
<211> 43  
<212> DNA  
<213> Artificial Sequence

<220>  
<221> primer\_bind  
<223> reverse primer useful for generation of  
PC4:HSA albumin fusion VECTOR

<220>  
<221> misc\_feature  
<222> (6)..(11)  
<223> Asp718 restriction site

<220>  
<221> misc\_feature  
<222> (12)..(17)  
<223> EcoRI restriction site

<220>  
<221> misc\_feature  
<222> (15)..(17)  
<223> reverse complement of stop codon

<220>  
<221> misc\_feature  
<222> (18)..(25)  
<223> AscI restriction site

<220>  
<221> misc\_feature  
<222> (18)..(43)  
<223> reverse complement of DNA sequence encoding last 9 amino acids

<400> 31

gcagcgggtac cgaattcggc ggcgccttata agcctaaggc agc

43

<210> 32

<211> 46

<212> DNA

<213> Artificial Sequence

<220>

<221> primer\_bind

<223> forward primer useful for inserting Therapeutic protein into pC4:HSA vector

<220>

<221> misc\_feature

<222> (29)

<223> n equals a,t,g, or c

<220>

<221> misc\_feature

<222> (30)

<223> n equals a,t,g, or c

<220>

<221> misc\_feature

<222> (31)

<223> n equals a,t,g, or c

<220>

<221> misc\_feature

<222> (32)

<223> n equals a,t,g, or c

<220>

<221> misc\_feature

<222> (33)

<223> n equals a,t,g, or c

<220>

<221> misc\_feature

<222> (34)

<223> n equals a,t,g, or c

<220>

<221> misc\_feature

<222> (35)

<223> n equals a,t,g, or c

<220>

<221> misc\_feature

<222> (36)

<223> n equals a,t,g, or c

<220>

<221> misc\_feature

<222> (37)

<223> n equals a,t,g, or c

<220>

<221> misc\_feature

<222> (38)

<223> n equals a,t,g, or c

<220>

<221> misc\_feature

<222> (39)

<223> n equals a,t,g, or c

<220>

<221> misc\_feature

<222> (40)

<223> n equals a,t,g, or c

<220>

<221> misc\_feature

<222> (41)

<223> n equals a,t,g, or c

<220>

<221> misc\_feature

<222> (42)

<223> n equals a,t,g, or c

<220>

<221> misc\_feature

<222> (43)

<223> n equals a,t,g, or c

<220>

<221> misc\_feature

<222> (44)

<223> n equals a,t,g, or c

<220>

<221> misc\_feature

<222> (45)

<223> n equals a,t,g, or c

<220>

<221> misc\_feature

<222> (46)

<223> n equals a,t,g, or c

<400> 32

ccgccgctcg aggggtgtgt ttctgtcgann nnnnnnnnnnn nnnnnnn

46

<210> 33

<211> 55

<212> DNA

<213> Artificial Sequence

<220>

<221> primer\_bind

<223> reverse primer useful for inserting Therapeutic protein into pC4:HSA vector

<220>

<221> misc\_feature

<222> (38)

<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (39)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (40)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (41)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (42)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (43)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (44)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (45)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (46)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (47)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (48)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (49)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (50)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (51)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (52)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (53)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (54)  
<223> n equals a,t,g, or c

<220>  
<221> misc\_feature  
<222> (55)  
<223> n equals a,t,g, or c

<400> 33  
agtcccatcg atgagcaacc tcactcttgt gtgcacnnnn nnnnnnnnnnn nnnnn 55

<210> 34  
<211> 17  
<212> PRT  
<213> Artificial Sequence

<220>  
<221> signal  
<223> Stanniocalcin signal peptide

<400> 34  
Met Leu Gln Asn Ser Ala Val Leu Leu Leu Val Ile Ser Ala Ser  
1 5 10 15

Ala

<210> 35  
<211> 22  
<212> PRT  
<213> Artificial Sequence

<220>  
<221> signal  
<223> Synthetic signal peptide

<400> 35  
Met Pro Thr Trp Ala Trp Trp Leu Phe Leu Val Leu Leu Ala Leu  
1 5 10 15

Trp Ala Pro Ala Arg Gly  
20

<210> 36

&lt;211&gt; 66

&lt;212&gt; PRT

&lt;213&gt; Agkistrodon piscivorus

&lt;400&gt; 36

Ile Thr Tyr Thr Asp Cys Thr Glu Ser Gly Gln Asn Leu Cys Leu Cys  
 1 5 10 15

Glu Gly Ser Asn Val Cys Gly Lys Gly Asn Lys Cys Ile Leu Gly Ser  
 20 25 30

Gln Gly Lys Asp Asn Gln Cys Val Thr Gly Glu Gly Thr Pro Lys Pro  
 35 40 45

Gln Ser His Asn Gln Gly Asp Phe Glu Pro Ile Pro Glu Asp Ala Tyr  
 50 55 60

Asp Glu  
 65

&lt;210&gt; 37

&lt;211&gt; 71

&lt;212&gt; PRT

&lt;213&gt; Agkistrodon piscivorus

&lt;400&gt; 37

Glu Ala Gly Glu Glu Cys Asp Cys Gly Ser Pro Glu Asn Pro Cys Cys  
 1 5 10 15

Asp Ala Ala Thr Cys Lys Leu Arg Pro Gly Ala Gln Cys Ala Glu Gly  
 20 25 30

Leu Cys Cys Asp Gln Cys Lys Phe Met Lys Glu Gly Thr Val Cys Arg  
 35 40 45

Ala Arg Gly Asp Asp Val Asn Asp Tyr Cys Asn Gly Ile Ser Ala Gly  
 50 55 60

Cys Pro Arg Asn Pro Phe His  
 65 70



# **INDICATIONS RELATING TO A DEPOSITED MICROORGANISM OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 58, line 12.

**B. IDENTIFICATION OF DEPOSIT**

Further deposits are identified on an additional sheet ☒

Name of depositary institution: American Type Culture Collection

Address of depositary institution (including postal code and country)

10801 University Boulevard  
Manassas, Virginia 20110-2209  
United States of America

Date of deposit

11 April 2001

Accession Number

PTA-3276

**C. ADDITIONAL INDICATIONS** (leave blank if not applicable)

This information is continued on an additional sheet ☐

**D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE** (if the indications are not for all designated States)

Europe

In respect of those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28(4) EPC).

Continued on additional sheets

**E. SEPARATE FURNISHING OF INDICATIONS** (leave blank if not applicable)

The indications listed below will be submitted to the international Bureau later (specify the general nature of the indications e.g. "Accession Number of Deposit")

	For receiving Office use only			For International Bureau use only	
<input type="checkbox"/> This sheet was received with the international application			<input checked="" type="checkbox"/> This sheet was received by the International Bureau on:		
			<div style="font-size: 1.2em; font-weight: bold;">16 MAY 2001</div>		
Authorized officer			Authorized officer		

**ATCC Deposit No.: PTA-3276**

## **CANADA**

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

## **NORWAY**

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

## **AUSTRALIA**

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

## **FINLAND**

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

## **UNITED KINGDOM**

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

**ATCC Deposit No.: PTA-3276**

## **DENMARK**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

## **SWEDEN**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

## **NETHERLANDS**

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

# **INDICATIONS RELATING TO A DEPOSITED MICROORGANISM OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

**A.** The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 58, line 12

**B. IDENTIFICATION OF DEPOSIT**

Further deposits are identified on an additional sheet ☒

Name of depositary institution: American Type Culture Collection

Address of depositary institution (including postal code and country)  
10801 University Boulevard  
Manassas, Virginia 20110-2209  
United States of America

Date of deposit

11 April 2001

Accession Number

PTA-3277

**C. ADDITIONAL INDICATIONS** (leave blank if not applicable)

This information is continued on an additional sheet ☐

**D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE** (if the indications are not for all designated States)


Europe

In respect of those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28(4) EPC).

Continued on additional sheets

**E. SEPARATE FURNISHING OF INDICATIONS** (leave blank if not applicable)

The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g. "Accession Number of Deposit")

	For receiving Office use only			For International Bureau use only	
<input type="checkbox"/> This sheet was received with the international application			<input checked="" type="checkbox"/> This sheet was received by the International Bureau on <div style="text-align: center;">16 MAY 2001</div>		
Authorized officer			Authorized officer 		

**ATCC Deposit No.: PTA-3277**

## **CANADA**

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

## **NORWAY**

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

## **AUSTRALIA**

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

## **FINLAND**

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

## **UNITED KINGDOM**

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

**ATCC Deposit No.: PTA-3277**

## **DENMARK**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

## **SWEDEN**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

## **NETHERLANDS**

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

**INDICATIONS RELATING TO A DEPOSITED MICROORGANISM  
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

**A.** The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 58, line 12.

**B. IDENTIFICATION OF DEPOSIT**

Further deposits are identified on an additional sheet ☒

Name of depositary institution: American Type Culture Collection

Address of depositary institution (including postal code and country)

10801 University Boulevard  
Manassas, Virginia 20110-2209  
United States of America

Date of deposit

11 April 2001

Accession Number

PTA-3278

**C. ADDITIONAL INDICATIONS** (leave blank if not applicable)

This information is continued on an additional sheet ☐

**D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE** (if the indications are not for all designated States)

Europe

In respect of those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28(4) EPC).

Continued on additional sheets

**E. SEPARATE FURNISHING OF INDICATIONS** (leave blank if not applicable)

The indications listed below will be submitted to the international Bureau later (specify the general nature of the indications e.g. "Accession Number of Deposit")

	For receiving Office use only			For International Bureau use only	
<input type="checkbox"/> This sheet was received with the international application			<input checked="" type="checkbox"/> This sheet was received by the International Bureau on:		
			16 MAY 2001		
Authorized officer			Authorized officer		

**ATCC Deposit No.: PTA-3278**

**CANADA**

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

**NORWAY**

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

**AUSTRALIA**

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

**FINLAND**

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

**UNITED KINGDOM**

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.



**ATCC Deposit No.: PTA-3278**

## **DENMARK**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

## **SWEDEN**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

## **NETHERLANDS**

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

# **INDICATIONS RELATING TO A DEPOSITED MICROORGANISM OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

**A.** The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 58, line 12.

**B. IDENTIFICATION OF DEPOSIT**

Further deposits are identified on an additional sheet ☒

Name of depositary institution: American Type Culture Collection

Address of depositary institution (including postal code and country)

10801 University Boulevard  
Manassas, Virginia 20110-2209  
United States of America

Date of deposit

11 April 2001

Accession Number

PTA-3279

**C. ADDITIONAL INDICATIONS** (leave blank if not applicable)

This information is continued on an additional sheet ☐

**D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE** (if the indications are not for all designated States)

Europe

In respect of those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28(4) EPC).

Continued on additional sheets

**E. SEPARATE FURNISHING OF INDICATIONS** (leave blank if not applicable)

The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")

	For receiving Office use only			For International Bureau use only	
<input type="checkbox"/> This sheet was received with the international application			<input checked="" type="checkbox"/> This sheet was received by the International Bureau on <div style="text-align: center;">16 MAY 2001</div>		
Authorized officer			Authorized officer		

**ATCC Deposit No.: PTA-3279**

## **CANADA**

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

## **NORWAY**

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

## **AUSTRALIA**

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

## **FINLAND**

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

## **UNITED KINGDOM**

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

**ATCC Deposit No.: PTA-3279****DENMARK**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

**SWEDEN**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

**NETHERLANDS**

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US01/12009

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(7) : C07K 14/00; C07H 21/04; C12N 15/00; A61K 38/16

US CL : 435/320.1; 514/2; 530/350; 536/23.4

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/320.1; 514/2; 530/350; 536/23.4

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

MEDLINE, BIOSIS, EMBASE, CAPLUS, various sequence databases

search terms: chimera, fusion, human serum albumin, interferon, cytokine, erythropoietin, SEQ ID No. 18

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X Y	WO 99/66054 A2 (GENZYME TRANSGENICS CORP.) 23 December 1999, see entire document.	1,2,4,6,9,10,12,14,16,17,24,28,30-37,40-42 3,15,25,26,29,39 N
X Y	WO 97/24445 A1 (DELTA BIOTECHNOLOGY LIMITED) 10 July 1997, see entire document, especially pp. 3-6, 12, 13, and 19-26	1-4,6,9,10,13-17,25,28-34,36,37,39-42 5,7,8,24,26,27,35,38

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

"	Special categories of cited documents:	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A"	document defining the general state of the art which is not considered to be of particular relevance	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"B"	earlier document published on or after the international filing date	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&"	document member of the same patent family
"O"	document referring to an oral disclosure, use, exhibition or other means		
"P"	document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search 16 JULY 2001	Date of mailing of the international search report 22 AUG 2001
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer TERRY J. DEY BRONWEN M. LOPEZ PARALEGAL SPECIALIST TECHNOLOGY CENTER 1600 Telephone No. (703) 305-3230

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US01/12009

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X <u>Y</u>	US 5,705,363 A (IMAKAWA, K.) 6 January 1998, see entire document, especially col. 23, line 44-col. 24, line 56.	1,2,4,5-7,10,2 5,33-38,40-42
X	US 5,766,883 A (BALLANCE et al) 16 June 1998, see entire document, especially col. 1, line 38-54, col. 6, line 53-col 10, line 41, col. 11, line 36-col. 12, line 40.	3,14,15,16,17,24, 1,2,4,6,10,13, 14,25,30- 34,36,37,40-42